

PROBING THE IGM/GALAXY CONNECTION V: ON THE ORIGIN OF Ly α AND O VI ABSORPTION AT $Z < 0.2$

J. XAVIER PROCHASKA¹, B. WEINER², H.-W. CHEN³, J. MULCHAEY⁴, K. COOKSEY⁵,

Draft version January 14, 2013

ABSTRACT

We analyze the association of galaxies to Ly α and O VI absorption, the most commonly detected transitions in the low- z intergalactic medium (IGM), in the fields of 14 quasars with $z_{\text{em}} = 0.06$ – 0.57 . Confirming previous studies, we observe a high covering fraction for Ly α absorption to impact parameter $\rho = 300 h_{72}^{-1}$ kpc: 33/37 of our $L > 0.01L^*$ galaxies show Ly α equivalent width $W^{\text{Ly}\alpha} \geq 50$ mÅ. Galaxies of all luminosity $L > 0.01L^*$ and spectral type are surrounded by a diffuse and ionized circumgalactic medium (CGM), whose baryonic mass is estimated at $\sim 10^{10.5 \pm 0.3} M_{\odot}$ for a constant N_{H} . The virialized halos and extended CGM of present-day galaxies are responsible for most strong Ly α absorbers ($W^{\text{Ly}\alpha} > 300$ mÅ) but cannot reproduce the majority of observed lines in the Ly α forest. We conclude that the majority of Ly α absorption with $W^{\text{Ly}\alpha} = 30$ – 300 mÅ occurs in the cosmic web predicted by cosmological simulations and estimate a characteristic width for these filaments of $\approx 400 h_{72}^{-1}$ kpc. Regarding O VI, we observe a near unity covering fraction to $\rho = 200 h_{72}^{-1}$ kpc for $L > 0.1L^*$ galaxies and to $\rho = 300 h_{72}^{-1}$ kpc for sub- L^* ($0.1L^* < L < L^*$) galaxies. Similar to our Ly α results, stronger O VI systems ($W^{1031} > 70$ mÅ) arise in the virialized halos of $L > 0.1L^*$ galaxies. Unlike Ly α , the weaker O VI systems ($W^{1031} \approx 30$ mÅ) arise in the extended CGM of sub- L^* galaxies. The majority of O VI gas observed in the low- z IGM is associated with a diffuse medium surrounding individual galaxies with $L \approx 0.3L^*$, and rarely originates in the so-called warm-hot IGM (WHIM) predicted by cosmological simulations.

Subject headings: absorption lines – intergalactic medium – Lyman limit systems

1. INTRODUCTION

Ultraviolet absorption-line spectroscopy remains the most efficient and sensitive means of studying the diffuse gas that permeates the universe. This gas, commonly referred to as the intergalactic medium (the IGM) or Ly α forest, is the dominant reservoir of baryons in the universe at all epochs (Prochaska & Tumlinson 2008). It provides fresh fuel for star-formation and collects the radiation and metals produced by galaxies and active galactic nuclei. Furthermore, the IGM is expected to trace the dark matter distribution of large-scale structure, offering unique constraints on our cosmological paradigm (e.g. Miralda-Escudé et al. 1996; Rauch 1998; McDonald et al. 2006; Viel et al. 2009).

Analysis of quasar spectra acquired with UV-sensitive spectrometers on the *Hubble Space Telescope* (*HST*) and *Far-Ultraviolet Spectroscopic Explorer* (*FUSE*) have surveyed the redshift distribution, metal-enrichment, ionization state, and temperature of the IGM in the $z < 1$ universe (e.g. Weymann et al. 1998; Davé et al. 1999;

Prochaska et al. 2006; Thom & Chen 2008b; Tripp et al. 2008; Danforth & Shull 2008; Cooksey et al. 2010). These studies have demonstrated, similar to the high z IGM, that the gas collects in a set of relatively discrete ‘lines’ in redshift, has a high ionization fraction, and is often enriched with heavy metals (e.g. C, O). The Ly α forest is significantly sparser at $z < 1$, however, because of the steady expansion of the universe. The cosmic abundance of heavy metals also appears to have evolved (e.g. Prochaska et al. 2004; Danforth & Shull 2008; Cooksey et al. 2010), although the scatter for individual regions is large.

Over the past decade, extra attention has been placed on the search for a warm-hot phase of the IGM ($T \gtrsim 10^6$ K) that is predicted by cosmological simulations to be a major baryonic reservoir in the low z universe (Cen & Ostriker 1999; Davé et al. 2001). In the UV, the search for this so-called WHIM (warm-hot intergalactic medium) includes surveys for the O VI doublet (e.g. Tripp et al. 2008; Thom & Chen 2008b) searches for thermally broadened Ly α lines (e.g. Lehner et al. 2007) and, most recently, surveys for Ne VIII absorption (Savage et al. 2005; Narayanan et al. 2009). Owing to its greater ease of detection, the O VI doublet has been the most frequently studied transition to date. Focused surveys have characterized its incidence as a function of equivalent width and column density (Tripp et al. 2008; Thom & Chen 2008b; Danforth & Shull 2008; Wakker & Savage 2009), and have permitted statistical comparison with predictions from cosmological simulations (e.g. Cen et al. 2009; Oppenheimer & Davé 2009).

¹ Department of Astronomy and Astrophysics & UCO/Lick Observatory, University of California, 1156 High Street, Santa Cruz, CA 95064; xavier@ucolick.org

² Steward Observatory, University of Arizona, 933 N. Cherry Ave., Tucson, AZ 85721; bjw@as.arizona.edu

³ Department of Astronomy; University of Chicago; 5640 S. Ellis Ave., Chicago, IL 60637; hchen@oddjob.uchicago.edu

⁴ Carnegie Observatories; 813 Santa Barbara St., Pasadena, CA 91101; mulchaey@obs.carnegiescience.edu

⁵ NSF Astronomy & Astrophysics Postdoctoral Fellow, MIT Kavli Institute for Astrophysics & Space Research, 77 Massachusetts Avenue, 37-611, Cambridge, MA 02139, USA; kcooksey@space.mit.edu

Although UV spectral analysis provides robust constraints on the nature and distribution of diffuse gas, considerable debate remains over the origin and cosmological relevance of the observed absorption systems. For example, hot and diffuse gas is predicted (and observed) to exist in the outer halos of individual galaxies (Spitzer 1956; Bahcall & Spitzer 1969; Mo & Miralda-Escude 1996; Maller & Bullock 2004; Sembach et al. 2003), the intra-group medium (e.g. Mulchaey et al. 1996; Freeland et al. 2008), and also the filamentary structures of the IGM (e.g. Cen & Ostriker 1999). The incidence of O VI, therefore, is likely a sensitive function of how metals are dispersed on galactic and intergalactic scales and also the physical conditions within these various environments (e.g. Oppenheimer & Davé 2009). Not surprisingly, groups have drawn competing conclusions on whether the O VI absorbers primarily trace the low-density WHIM or a photoionized, cooler phase. Similarly, multiple scenarios may explain the incidence and properties of the Ly α ‘clouds’ along single IGM sightlines. While ram-pressure stripping, tidal stripping, and galactic-scale winds are all believed to play a role in transporting gas from galaxies to the IGM, the timing and relative importance of each of these remain open issues. In short, there remain several fundamental questions on the origin and nature of the two most common transitions of the IGM.

Researchers have long recognized that additional insight into the IGM may be gained by surveying the fields surrounding absorption-line systems for galaxies and their large-scale structures. Early efforts focused on Ly α absorption and the relation of HI gas to galaxies and large-scale structures (Spinrad et al. 1993; Morris et al. 1993; Lanzetta et al. 1995). A key result was that sightlines with small impact parameters to galaxies ($\rho \lesssim 250 h_{72}^{-1}$ kpc) had a high incidence of moderate to strong Ly α absorption (Lanzetta et al. 1995; Chen et al. 1998; Tripp et al. 1998; Chen et al. 2001b; Bowen et al. 2002; Morris & Jannuzi 2006). This indicates galaxies are surrounded by a diffuse, highly ionized medium that gives rise to significant Ly α absorption (see also Wakker & Savage 2009). These studies were followed by two-point cross-correlation analysis of galaxies to absorbers on ≈ 1 Mpc scales (Chen et al. 2005; Wilman et al. 2007; Chen & Mulchaey 2009) and have recently been extended to $z \sim 1$ (Shone et al. 2010). Their results reveal significant clustering between galaxies and strong Ly α systems (absorbers with HI column density, $N_{\text{HI}} > 10^{15} \text{ cm}^{-2}$) that implies a causal connection. In contrast, they measure a very weak or no clustering signal for low- N_{HI} absorption systems which implies the two phenomena trace different structures in the universe. While previous work has offered valuable insight into the nature of the Ly α forest, it is limited by sample variance both in terms of the number of absorbers, galaxies, and fields surveyed.

Several projects have now also considered the galaxy/absorber connection for the metal-enriched IGM, e.g. Mg II, C IV, and O VI selected systems. For the first, observers have commonly associated strong Mg II absorption with individual galaxies at impact parameters $\rho < 100$ kpc, by searching for galaxies at the redshifts of known Mg II systems (e.g. Bergeron 1986;

Bergeron & Boisse 1991). Indeed, galaxies at low- z show modest to high covering fractions of cool gas ($T \sim 10^4$ K) traced by Mg II absorption (Barton & Cooke 2009; Chen & Tinker 2008; Chen et al. 2010). This has led to the association of metal-enriched, cool IGM gas with the ‘halos’ of individual galaxies. Galaxies also exhibit a high covering fraction to C IV absorption for impact parameters $\rho \lesssim 200$ kpc (Chen et al. 2001a). Although the origin of this highly ionized gas is not well established, these results likely require a multi-phase medium in the virialized halos of modern galaxies (Mo & Miralda-Escude 1996).

The O VI doublet has long been recognized to trace a highly ionized, warm/hot phase of the Galaxy. The original detections associated O VI gas with a Local Bubble surrounding the Sun (York 1974; Savage & Lehner 2006). A statistical analysis of O VI absorption towards extragalactic sources with *FUSE* suggests a thick layer of highly ionized gas with a scale height of ≈ 3 kpc (Savage et al. 2000). Lastly, studies of O VI absorption toward extragalactic sources reveal a high covering fraction from an inferred halo of collisionally ionized gas at $T \approx 10^5$ K (Savage & de Boer 1979; Savage et al. 2003; Sembach et al. 2006). Similar halos of O VI-bearing gas have been inferred for external galaxies through O VI emission maps (Bregman et al. 2005). It is evident that some fraction of O VI observed in the IGM must arise from galactic halos.

Connecting O VI gas of the IGM to galaxies and the structures they reside within has been the focus of several recent studies. In Prochaska et al. (2006), we examined the field surrounding the multiple O VI systems identified along the sightline to PKS0405–123. While the strong absorption at $z = 0.167$ was notable for its association with two galaxies at small impact parameter ($\rho \sim 100$ kpc), additional O VI systems showed no obvious galactic counterparts and/or galaxies only at large impact parameter. A similar diversity of galactic environment has been observed for other sightlines (Tumlinson et al. 2005; Sembach et al. 2004; Cooksey et al. 2008; Chen & Mulchaey 2009). Stocke et al. (2006) performed the first multi-sightline analysis on the galaxy/absorber connection for O VI. For the 9 O VI systems in their fields complete to $0.1 L^*$ galaxies, they estimated a median distance to an $L \geq 0.1 L^*$ galaxy of $\approx 180 h_{70}^{-1}$ kpc. Furthermore, they identified an L^* galaxy within $800 h_{70}^{-1}$ kpc for essentially each of the 23 O VI absorbers in their survey. They inferred, based on these results, that O VI gas does not occur in galaxy voids. They further hypothesized that galaxies with $L \leq 0.1 L^*$ galaxies may be most responsible for O VI absorption. Similar results were found by Wakker & Savage (2009) for a small sample of $z \approx 0$ galaxies.

Recently, we published the results of a galaxy survey performed with the WFCCD spectrometer on the 100" Dupont telescope at Las Campanas Observatory (LCO) in the fields surrounding 20 UV-bright quasars at $z < 1$ (Prochaska et al. 2011). The principal motivation for this survey was to assess the galaxy/absorber connection for O VI gas at $z \leq 0.2$. In this paper, we present the first scientific results from our complete survey. Previous papers studied the galaxy/IGM connection in a few, indi-

TABLE 1
LCO/WFCCD FIELDS ANALYZED

Quasar	RA (J2000)	DEC (J2000)	z_{em}	<i>HST</i> UV Spectroscopic Datasets	<i>FUSE</i> ^a	Ly α Ref.	O VI Ref.	N_{spec}^b
Q0026+1259	00:29:13.8	+13:16:04	0.142	GHRs/(G270M)	20		9	60
TonS 180	00:57:20.0	-22:22:56	0.062	STIS/(G140M,G230MB)	132	5	4	7
PKS0312-77	03:11:55.2	-76:51:51	0.223	STIS/(E140M)		3	1,3	56
PKS0405-123	04:07:48.4	-12:11:37	0.573	STIS/(E140M,G230M); GHRs/(G160M,G200M)	71	3,4,9	1,3,4,9	565
PG1004+130	10:07:26.1	+12:48:56	0.240	STIS/(G140M)	85	9	9	61
HE1029-140	10:31:54.3	-14:16:51	0.086	STIS/(G140M)		5		8
PG1116+215	11:19:08.70	+21:19:18.	0.176	STIS/(G140M,E140M,E230M); GHRs/(G140L)	76	1,3,4,5,9	1,2,3,4	74
PG1211+143	12:14:17.7	+14:03:13.	0.081	STIS/(G140M,E140M); GHRs/(G140L,G270M)	52	3,4,5	3,4	25
PG1216+069	12:19:20.9	+06:38:38	0.331	STIS/(E140M); GHRs/(G140L)	13	1,3,6,9	1,2,3,9	101
3C273	12:29:6.70	+02:03:9.0	0.158	STIS/(E140M); GHRs/(FG130,FG190,G160M)	42	1,3,4,9	1,2,3,4	32
PKS1302-102	13:05:33.0	-10:33:19	0.286	STIS/(E140M)	140	3,4,9	1,3,4,9	63
MRK1383	14:29:06.4	+01:17:06.0	0.086	STIS/(E140M,G140M)	64	3,4,5	4,9	5
FJ2155-0922	21:55:01.5	-09:22:25.0	0.192	STIS/(E140M,G230MB)	46	1,3,4,9	1-4,9	105
PKS2155-304	21:58:51.8	-30:13:30.0	0.116	STIS/(E140M); GHRs/(G160M,ECH-B,G140L)	123	3,4,9	3,4	43

REFERENCES. — 1: Tripp et al. (2008); 2: Thom & Chen (2008b); 3: Danforth & Shull (2008); 4: Danforth et al. (2006); 5: Penton et al. (2004); 6: Chen & Mulchaey (2009); 9: This paper.

^a Total integration time in ks.

^b Number of spectroscopically determined galaxy redshifts for objects with $0.005 < z < z_{\text{em}}$.

vidual fields (Prochaska et al. 2004; Cooksey et al. 2008; Chen et al. 2005; Lehner et al. 2009). A future paper will study the cross-correlation function between galaxies and Ly α /O VI absorbers.

This paper is organized as follows. In § 2, we summarize the datasets that comprise the galaxy surveys (Prochaska et al. 2011) and the characterization of the IGM along the sightlines. The primary results of our analysis of galaxy/absorber association for Ly α and O VI are given in § 3. We discuss the implications for the low- z IGM of these results and their context with respect to previous work in § 4 and we conclude with a summary in § 5. Unless otherwise specified, we adopt a $\Omega_{\Lambda} = 0.72$, $\Omega_m = 0.28$, $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$ cosmology (Dunkley et al. 2009). Furthermore, all distances are quoted in proper units unless otherwise noted.

2. DATASETS

2.1. The LOC/WFCCD Galaxy Survey

We have recently published a survey of galaxies in the fields of 20 quasars with bright UV fluxes (Paper IV), the majority of which have UV spectral datasets that yield precise measurements of Ly α and/or O VI absorption (Table 1). The galaxy survey was performed with the WFCCD spectrometer on the 100" Dupont telescope and was designed to achieve a high level of completeness to $R \leq 19.5 \text{ mag}$ galaxies in an $\approx 20' \times 20'$ field centered on each quasar. These survey parameters were chosen to recover dwarf galaxies ($L < 0.1 L^*$) to at least $z = 0.1$ and to span a $\approx 1 \text{ Mpc}$ impact parameter at $z \approx 0.1$. With this experimental design, we aimed to study the galaxies and their structures (e.g. groups) associated with IGM absorption. In several analyses (where noted), we also include the galaxy survey in the field of PKS0405-123 published by Williger et al. (2006).

The fields were selected on the basis of the quasar redshift (higher z was preferred) and UV flux, with some preference given to fields with existing UV spectroscopic datasets. None of the fields were chosen because of the presence of known absorption systems, although an O VI-bearing Lyman limit system at $z = 0.167$ toward

PKS0405-123 had been analyzed by our group previously (Chen & Prochaska 2000). As such, the galaxies discovered by our survey provide an unbiased sample that can then be searched for associated hydrogen and metal-line absorption.

This LCO/WFCCD galaxy survey provides the galaxy sample for the following analysis; we consider the results from complementary surveys in the discussion that follows (§ 4). Altogether, the LCO/WFCCD galaxy survey comprises 1198 galaxies with $0.005 < z < (z_{\text{em}} - 0.01)$ with a median redshift of 0.18. The distribution of their luminosities peaks near $\approx 0.3 L^*$ and extends to $0.01 L^*$ and $\approx 5 L^*$.

2.2. IGM Surveys for Ly α and O VI Absorption

In the following IGM/galaxy analysis, we leverage published surveys for Ly α and O VI absorption in the quasar spectra of our fields. This corresponds to 13 sightlines with UV spectra from the *HST*/GHRs, *HST*/STIS, and/or *FUSE* instruments (Table 1). We restrict our analysis to surveys based on high spectral resolution data which generally provide equivalent width sensitivity $\sigma(W)$ to a few tens of mÅ. Although the published surveys occasionally disagree on the identification of a particular line or on the confidence of its detection, we have not attempted to reconcile these arguments. Instead, we adopt all published detections as bona-fide systems. When multiple groups have reported measurements for the same system, we have adopted the values from publications in this order: [Ly α] – Tripp et al. (2008); Danforth & Shull (2008); Chen & Mulchaey (2009); Danforth et al. (2006); Penton et al. (2004); [O VI] – Tripp et al. (2008); Thom & Chen (2008b); Danforth & Shull (2008); Danforth et al. (2006). Because we rely primarily on published surveys (see below), the IGM results were derived independently of our galaxy survey.

For galaxies within several hundred kpc (up to 1 Mpc for Ly α) of these sightlines, we have also done a focused search for associated Ly α and O VI absorption at their redshifts. Specifically, if there were no published mea-

TABLE 2
NEW TARGETED SEARCH FOR $\text{Ly}\alpha$ AND O VI ABSORPTION ASSOCIATED WITH GALAXIES

Quasar	RA (J2000)	DEC (J2000)	z_{gal}	$W^{\text{Ly}\alpha}$ (Å)	N_{HI}	W^{OVI} (mÅ)	N_{OVI}	UV Spectra
Q0026+1259	00:29:13.8	+13:16:04	0.03290	...	...	< 120	< 14.0	FUSE
			0.03930	...	...	< 120	< 14.0	FUSE
			0.09610	...	...	< 100	< 14.0	FUSE
			0.11250	...	...	< 100	< 14.0	FUSE
PKS0405-123	04:07:48.4	-12:11:37	0.04320	< 0.03	< 12.7	...	...	STIS/E140M
			0.07910	< 0.03	< 12.7	< 40	< 13.5	FUSE, STIS/E140M
			0.08000	< 0.03	< 12.7	< 40	< 13.5	FUSE, STIS/E140M
			0.15220	...	...	< 50	< 13.6	STIS/E140M
			0.20300	< 0.05	< 13.0	< 30	< 13.4	STIS/E140M
			0.20890	< 0.03	< 12.7	...	...	STIS/E140M
			0.24840	< 0.05	< 12.9	< 30	< 13.4	STIS/E140M
			0.28000	< 0.03	< 12.8	...	...	STIS/E140M
			0.29510	< 0.09	< 13.2	< 30	< 13.4	STIS/E140M
			0.29760	...	...	< 50	< 13.6	STIS/E140M
			0.30990	< 0.04	< 12.9	< 40	< 13.5	STIS/E140M
			0.32030	< 0.05	< 13.0	< 30	< 13.4	STIS/E140M
			0.32520	...	...	< 50	< 13.6	STIS/E140M
			0.34150	< 0.15	< 13.5	...	...	STIS/E140M
			0.36810	< 0.05	< 13.0	...	...	STIS/E140M
			0.37850	< 0.06	< 13.0	...	...	STIS/E140M
PG1004+130	10:07:26.1	+12:48:56	0.00920	0.26 ± 0.03	13.9 ± 0.15	< 50	< 13.6	FUSE, STIS/G140M
			0.03008	...	...	< 200	< 14.5	FUSE
			0.07040	...	...	< 100	< 13.9	FUSE
			0.16740	...	...	< 80	< 13.8	STIS/G140M
			0.19310	...	...	< 80	< 13.8	STIS/G140M
PG1216+069	12:19:20.9	+06:38:38	0.00630	2.32 ± 0.06	19.3 ± 0.05	< 200	< 14.2	FUSE, STIS/E140M
			0.00810	< 0.06	< 13.1	< 200	< 14.2	FUSE, STIS/E140M
			0.01262	0.35 ± 0.03	> 14.1	< 200	< 14.2	FUSE, STIS/E140M
			0.08150	...	...	< 120	< 14.0	FUSE
			0.11840	< 0.03	< 12.7	< 130	< 14.1	FUSE, STIS/E140M
			0.11910	< 0.03	< 12.7	< 130	< 14.1	FUSE, STIS/E140M
			0.13540	...	...	< 130	< 14.1	FUSE
			0.18100	...	...	< 80	< 13.8	STIS/E140M
			0.19170	< 0.05	< 12.9	< 40	< 13.5	STIS/E140M
			0.24640	< 0.05	< 12.9	< 40	< 13.5	STIS/E140M
			0.27950	< 0.05	< 12.9	< 60	< 13.6	STIS/E140M
PKS1302-102	13:05:33.0	-10:33:19	0.28000	< 0.05	< 12.9	< 60	< 13.6	STIS/E140M
			0.02520	< 0.10	< 13.3	< 40	< 13.5	FUSE, STIS/E140M
			0.03650	< 0.25	< 15.0	...	...	STIS/E140M
			0.05690	< 0.35	< 15.0	< 40	< 13.5	FUSE, STIS/E140M
			0.07120	< 0.50	< 16.0	< 40	< 13.5	FUSE, STIS/E140M
			0.07180	< 0.50	< 16.0	< 40	< 13.5	FUSE, STIS/E140M
			0.10620	< 0.05	< 12.9	...	...	STIS/E140M
			0.13930	< 0.04	< 12.8	...	...	STIS/E140M
MRK1383	14:29:06.4	+01:17:06.0	0.14290	< 0.04	< 12.9	< 40	< 13.5	FUSE, STIS/E140M
			0.02990	...	...	< 40	< 13.5	FUSE

NOTE. — The N_{HI} value for the $z = 0.0063$ absorber toward PG1216+069 is taken from Tripp et al. (2006)

surement near the redshift of the galaxy yet spectra exist we reanalyzed the data ourselves. Furthermore, sufficient quality UV spectra exist for $\text{Ly}\alpha$ and/or O VI analysis in one of the LCO/WFCCD survey fields yet none has been published (Q0026+1259). We have also done a focused analysis in this sightline according to galaxies discovered in the LCO/WFCCD survey. To perform this new IGM analysis, we used spectra reduced by the instrument pipelines as described in Cooksey et al. (2008). We normalized the quasar continuum with automated algorithms and measured equivalent widths (and errors) with simple boxcar summation. If no absorption was present within $\pm 400 \text{ km s}^{-1}$ of the galaxy redshift, we report a 3σ upper limit to the equivalent width and column density (assuming the linear curve-of-growth approximation). For positive detections, we measured column densities with the apparent optical depth method (Savage & Sembach 1991). In a few cases, the predicted

wavelength of a $\text{Ly}\alpha$ or O VI line coincides with Galactic ISM absorption or a coincident transition from another absorption system. In these events, we adopt the equivalent width of the coincident line as an upper limit to the equivalent width.

The measurements for our IGM analysis are summarized in Table 2. Altogether, we have useful constraints on $\text{Ly}\alpha$ and/or O VI absorption for 14 of our WFCCD survey fields.

3. RESULTS

We take two complementary approaches to the analysis: (i) we examine all galaxies that lie close to the sightlines for associated $\text{Ly}\alpha$ and O VI absorption; and (ii) we search the fields for galaxies associated with detected $\text{Ly}\alpha$ and O VI absorbers. Each of these techniques has specific strengths and weaknesses for interpreting the IGM/galaxy connection, as described below. Together,

however, they provide powerful insight into the nature of the IGM and its interplay with galaxies at $z \sim 0$.

In the following, we establish the velocity criterion for associating a galaxy with IGM absorption (or vice-versa) to be a velocity difference $|\delta v| < 400 \text{ km s}^{-1}$. This value was chosen to be large enough to account for error in the galaxy redshifts (Prochaska et al. 2011) and to allow for velocities between the galaxy and gas that are characteristic of galactic dynamics. We emphasize that over 80% of the purported galaxy/absorber associations have $|\delta v| \leq 250 \text{ km s}^{-1}$ and that our results would not change qualitatively if we adopted a value lower than 400 km s^{-1} .

We examine the results with particular emphasis on the galaxy luminosity. These were derived from the apparent R -band magnitude, redshift, and spectral type (see Prochaska et al. 2011). Generally, we consider three luminosity intervals: (1) $L < 0.1L^*$, termed dwarf galaxies; (2) $0.1L^* < L < L^*$, termed sub- L^* galaxies and; (3) $L > L^*$, termed L^* galaxies. For these definitions, we adopt the r -band value for L^* as measured from the Sloan Digital Sky Survey which corresponds to $z \approx 0.1$: $r^* = -20.44 + 5 \log h_{100}$ (Blanton et al. 2003). With the cosmology we have adopted, this yields $r^* = -21.12 \text{ mag}$. We are motivated to this approach by the expectation that galaxies of a similar luminosity will, on average, trace similar environments and have dark matter halos with common characteristics. Ideally, we might separate the galaxies by their dark matter halo mass, as inferred from the galaxy’s kinematics. Unfortunately, our spectra are of insufficient quality to discriminate on kinematics nor do we have the necessary photometry to estimate stellar masses. Regarding the galaxy spectral-type, we do make the distinction between early-type (absorption-line dominated) and late-type (emission-line dominated) galaxies based on our spectra (see Prochaska et al. 2011, for more details).

When discussing the results, we make a crude distinction between the virialized halo of a galaxy defined by a virial radius r_{vir} and the circumgalactic medium (CGM) that surrounds it to a larger radius r_{CGM} . The motivation for this distinction is that galactic-scale processes (e.g. accretion, shock-heating, outflows) may be especially active within the virialized halo, and therefore produce a medium that is qualitatively different from the surrounding CGM. Furthermore gas at $r < r_{\text{vir}}$ may have a much greater probability of being gravitationally bound to the galaxy. At the same time, we doubt that r_{vir} marks a sharp boundary of any sort. Lastly, we note that a sightline which penetrates a galactic halo may show absorption from the gas within it and the gas that surrounds it.

In standard numerical and analytic analysis, r_{vir} is predicted to scale with the dark matter halo mass as $r_{\text{vir}} \propto M_h^{1/3}$ and with the circular velocity as $r_{\text{vir}} \propto v_c$. If we were to adopt the observed R -band Tully-Fisher (TF) relation $L \propto v_c^{3.4}$ (Courteau et al. 2007), we would infer that r_{vir} scales as $r_{\text{vir}} \propto L^{0.3}$. More modern treatments compare the clustering of galaxies as a function of luminosity to predictions for dark matter halos and thereby derive a relation between L and M_h (e.g. Zheng et al. 2007; The SDSS Collaboration et al. 2010). These results, which are dominated by $L > L^*$ galaxies, imply $M_h \propto L^{3.8}$ (in R -band) at high luminosity and a much

flatter relation at low luminosity (consistent with the higher M/L ratio inferred for low-mass galaxies). Therefore, while most treatments — empirical and theoretical — tend to agree that $r_{\text{vir}} \approx 250\text{--}300 \text{ kpc}$ for an L^* galaxy, various prescriptions yield very different estimates for the virial radius of low luminosity galaxies. In the following, we adopt characteristic values of $r_{\text{vir}} \approx 100 \text{ kpc}$ for the dwarf galaxies, $r_{\text{vir}} \approx 160 \text{ kpc}$ for a sub- L^* galaxy, and $r_{\text{vir}} \approx 250 \text{ kpc}$ for an L^* galaxy. This corresponds to a scaling relation:

$$r_{\text{vir}} = r_{\text{vir}}^* \left(\frac{L}{L^*} \right)^\beta \quad (1)$$

with $r_{\text{vir}}^* = 250 \text{ kpc}$ and $\beta = 0.2$. These assignments and this relation should be considered crude estimates and primarily serve to guide the discussion.

3.1. Gas Associated with Galaxies at Low Impact Parameter

In this subsection, we address the following question: granted that a galaxy with a certain luminosity lies at a given impact parameter from a quasar sightline, what is the probability of detecting Ly α or O VI absorption at that same redshift (within a given velocity interval δv) to a given equivalent width limit? More generally, we consider the distribution of equivalent widths and ionic column densities as a function of the galaxy luminosity L and impact parameter ρ . In principle, such measurements reveal the physical conditions of gas in the virialized halos of galaxies and in the circumgalactic medium (CGM) that surrounds them (e.g. filaments, the intragroup medium). In turn, the measurements may inform whether the gas is physically connected to the galaxy (e.g. gravitationally bound to its dark matter halo) and/or which galaxies may ‘produce’ IGM absorption at specific equivalent widths and column density.

There is (at least) one significant challenge to this analysis: galaxies cluster with one another. Therefore, if a galaxy lies at close impact parameter to a sightline there is a significant probability that one or more additional galaxies will also lie close to the same sightline. If we then detect IGM absorption at that redshift, there is an ambiguous association between the gas and each galaxy. One may be able to design an experiment that studied only isolated galaxies, but this would likely be unrepresentative of the full galaxy population. This issue is further complicated by the imperfections of our LCO/WFCCD survey. These include (i) the magnitude limit which implies sensitivity to fainter galaxies only at lower redshifts; (ii) the limited field-of-view which precludes a search for galaxies with $\rho \lesssim 100 \text{ kpc}$ at the lowest redshifts; and (iii) the varying sensitivity of the UV spectrometers with wavelength which dictates the equivalent width limit to Ly α and O VI absorption as a function of redshift. In the following, we emphasize these limitations where appropriate.

To (partially) address the ambiguity of multiple galaxies at a given redshift,⁶ we perform the analysis sepa-

⁶ Previous authors have introduced criteria based on galaxy clustering to select the galaxy of interest when two or more with similar redshift are located near the sightline (Chen et al. 2001b). In general, this amounts to selecting the galaxy with smallest impact parameter (Morris & Jannuzi 2006).




FIG. 1.— The four panels show the measured equivalent widths and column densities of Ly α and O VI absorption for all of the dwarf galaxies ($L < 0.1L^*$) in our LCO/WFCCD survey that lie within 300 kpc of a quasar sightline (where high quality UV spectroscopy exists). In each case, we report the maximum value for all Ly α (or O VI) absorption lines detected within 400 km s^{-1} of the galaxy redshift. Asterisks denote galaxies with late-type spectra and open circles signs denote early-type spectra (small/large arrows indicated limits for late/early-type galaxies). In cases where there is more than one dwarf galaxy within 400 km s^{-1} of one another, we plot only the one closest to the sightline. Upper limits denote non-detections or, in a few cases, unresolved blends with coincident absorption lines. The gray boxes (diamonds) denote systems where an additional sub- L^* (L^*) galaxy lies within 150 kpc of the sightline. Remarkably, these include nearly all of the systems associated with strong Ly α absorption and positive O VI detections. The gray triangles, meanwhile, denote galaxies associated with Virgo. Note the preponderance of Ly α associations but the rare detection of O VI gas, especially if one ignores the dwarf galaxies with a brighter neighbor near the sightline.

rately for dwarf, sub- L^* , and L^* galaxies. Initially, we examine all galaxies at impact parameter $\rho < 300 \text{ kpc}$ from the sightlines. This distance was arbitrarily chosen to exceed the virial radius of the galaxies yet to be small enough to primarily focus the analysis on individual galaxies. When two or more galaxies are separated by $|\delta v| < 400 \text{ km s}^{-1}$ and have similar luminosity, we only present the one closest to the sightline (this occurs very rarely).

Dwarf Galaxies: Consider first the $L < 0.1L^*$ galaxies (a.k.a. the dwarf galaxies); Table 3 lists the sample of systems at low impact parameter from our WFCCD survey. The table also lists the strongest Ly α and O VI absorbers within $|\delta v| \leq 400 \text{ km s}^{-1}$ of each galaxy, or an upper limit to the equivalent width and column density for non-detections. We also tabulate the closest galaxy with $L > 0.1L^*$ if one (or more) lies within 300 km s^{-1} of the dwarf galaxy’s redshift and 200 kpc of the sightline.

Figure 1 summarizes the absorption characteristics of the sample as a function of impact parameter. Regarding H I absorption, we note the positive detection

of Ly α absorption for 16/19 (84%) of the galaxies to $W^{\text{Ly}\alpha} = 100 \text{ mÅ}$ or $N_{\text{HI}} = 10^{13.5} \text{ cm}^{-2}$. We emphasize that unlike the high- z universe where the Ly α forest is sufficiently ‘dense’ that one identifies a line every $\sim 100 \text{ km s}^{-1}$, the probability of randomly associating a galaxy with an absorber is small at $z \sim 0$. Taking PG1116+215 as a representative sightline, the Ly α lines reported by Danforth & Shull (2008) cover $\approx 33\%$ of the sightline for $|\delta v| < 400 \text{ km s}^{-1}$ to $N_{\text{HI}} \approx 10^{13} \text{ cm}^{-2}$. Restricting to Ly α lines with $N_{\text{HI}} \geq 10^{13.5} \text{ cm}^{-2}$ (Figure 1), which is more characteristic of the gas associated with the dwarf galaxies, we find only $\approx 10\%$ of the sightline is covered. As such, if we randomly assigned redshifts to the galaxies with values uniformly sampling $z = [0, z_{\text{em}}]$, the probability of recovering 16/19 associations with Ly α lines having $N_{\text{HI}} > 10^{13.5} \text{ cm}^{-2}$ is negligible. We conclude, therefore, that Ly α absorption traces dwarf galaxies to impact parameters of 300 kpc or greater with a nearly unit covering fraction to $W^{\text{Ly}\alpha} = 100 \text{ mÅ}$ ($N_{\text{HI}} = 10^{13.5} \text{ cm}^{-2}$).

TABLE 3
DWARF GALAXIES WITHIN 300 KPC OF A QSO SIGHTLINE

Field	RA (J2000)	DEC (J2000)	z_{gal}	T ^a	ρ (kpc)	L (L^*)	$z_{\text{abs}}^{\text{Ly}\alpha}$	$W^{\text{Ly}\alpha}$ (mÅ)	N_{HI}	Rf	$z_{\text{abs}}^{\text{OVI}}$	W^{OVI} (mÅ)	N_{OVI}	Rf	ρ_n^b	L_n^b (L^*)
Q0026+1259	00:29:09.3	+13:16:29	0.0329	L	43	0.018	...	...	...	...	0.0329	< 120	< 14.0	9	...	...
TONS180	00:57:04.0	-22:26:51	0.0234	L	147	0.019	0.0234	222	13.80	5	0.0234	43	13.48	4	...	...
PKS0405-123	04:07:48.3	-12:11:02	0.1670	L	93	0.080	0.1671	697	15.47	3	0.1669	412	14.64	1	115	2.14
PG1004+130	10:07:06.5	+12:53:52	0.0092	L	73	0.007	0.0092	260	13.90	9	0.0092	< 50	< 13.6	9	...	...
	10:07:30.8	+12:53:50	0.0297	L	167	0.051	...	...	...	...	0.0301	< 200	< 14.50	9	...	...
HE1029-140	10:31:41.4	-14:12:49	0.0508	E	286	0.033	0.0516	278	14.00	5	...	...	...	...	...	...
PG1116+215	11:19:03.1	+21:16:24	0.0594	E	207	0.024	0.0593	187	13.64	1	0.0593	63	13.77	1	124	0.10
PG1211+143	12:14:06.9	+14:04:38	0.0520	L	172	0.089	0.0510	703	15.67	3	0.0513	...	14.30	4	131	0.83
	12:14:13.9	+14:03:31	0.0646	L	68	0.084	0.0645	535	15.73	3	0.0645	144	14.16	3	141	0.74
PG1216+069	12:18:38.7	+06:42:29	0.0067	E	86	0.029	0.0063	2320	19.30	9	0.0063	< 200	< 14.2	9	...	...
	12:19:20.7	+06:42:19	0.0081	E	34	0.008	0.0081	< 60	< 13.1	9	0.0081	< 200	< 14.2	9	...	...
	12:19:03.8	+06:33:43	0.0132	E	98	0.014	0.0126	288	13.94	6	0.0126	< 200	< 14.2	9	...	...
	12:19:14.5	+06:35:33	0.0805	E	294	0.061	0.0805	379	13.87	3	0.0815	< 120	< 14.0	9	...	...
3C273	12:29:50.6	+02:01:54	0.0062	E	79	0.006	0.0053	365	15.38	3	0.0053	< 15	< 13.2	4	...	...
PKS1302-102	13:05:58.0	-10:24:50	0.0252	L	299	0.009	0.0252	< 100	< 13.3	9	0.0252	< 40	< 13.5	9	...	...
MRK1383	14:29:42.4	+01:17:51	0.0281	E	285	0.009	0.0282	225	13.67	3	0.0283	< 16	< 13.1	4	...	...
	14:28:58.4	+01:13:06	0.0299	L	150	0.023	0.0299	66	13.10	5	0.0299	< 40	< 13.5	9	...	...
FJ2155-0922	21:54:56.8	-09:27:24	0.0504	E	282	0.045	0.0515	272	14.08	3	0.0501	< 68	< 13.9	4	...	...
	21:54:47.4	-09:23:05	0.0779	L	296	0.053	0.0777	448	> 14.3	1	0.0777	46	13.61	1	...	...
	21:54:52.3	-09:24:38	0.0808	E	272	0.070	0.0808	1007	15.11	3	0.0807	< 14	< 13.1	4	33	0.87
PKS2155-304	21:58:30.5	-30:11:02	0.0169	E	113	0.002	0.0170	143	13.56	3	0.0167	< 15	< 13.0	4	...	...

REFERENCES. — 1: Tripp et al. (2008); 2: Thom & Chen (2008b); 3: Danforth & Shull (2008); 4: Danforth et al. (2006); 5: Penton et al. (2004); 6: Chen & Mulchaey (2009); 9: This paper.

NOTE. — Upper limits correspond to 2σ constraints.

^a Spectral type (E=Early-type; L=Late-type; see text for the quantitative definition).

^b Impact parameter and luminosity of the brightest galaxy ($L > 0.1L^*$) within 200 kpc and $|\delta v| < 400 \text{ km s}^{-1}$ (if any).

Inspecting Figure 1 we note that the majority of dwarf galaxy-Ly α associations are at an impact parameter that is well beyond the presumed virial radius of $r_{\text{vir}} \approx 100 \text{ kpc}$. We infer, therefore, that the connection between these galaxies and H I absorption is not driven (generally) by galactic-scale phenomena, e.g. tidal debris, accreting gas, outflows. H I gas also arises in the extended CGM of these galaxies. The direct implication is that the two phenomena (Ly α absorption and dwarf galaxies) are simply highly correlated tracers of the same large-scale overdensity in the universe (e.g. a filament). This assertion is further supported by the fact that there is no strong correlation between the impact parameter and the Ly α absorption (in column density or equivalent width).

It is also evident from Figure 1 that a subset of the dwarf galaxies are associated with relatively strong Ly α absorbers, $W^{\text{Ly}\alpha} > 0.5 \text{ \AA}$ and $N_{\text{HI}} > 10^{15} \text{ cm}^{-2}$. However, most of these strong absorbers may also be associated to a brighter galaxy ($L > 0.1L^*$) within 200 kpc of the sightline. We have overplotted on the points gray boxes (diamonds) to indicate dwarf galaxies where an additional sub- L^* (L^*) galaxy lies within 150 kpc of the sightline. With the exception of the two galaxies associated with Virgo (at $z \approx 0.006$ toward PG1216+069 and 3C273; marked with gray triangles), all of the dwarf galaxies with associated strong Ly α absorption also have a brighter galaxy near the sightline. Furthermore, there are no cases where a brighter galaxy is nearby and the Ly α equivalent width is low. Although limited by small number statistics, these results suggest that the strongest H I absorbers are preferentially associated with a $L > 0.1L^*$ galaxy instead of the dwarf.

Lastly, with regards to dwarf galaxies and H I absorp-

tion, we consider the two non-detections in Ly α . One case is located at nearly 300 kpc from PKS1302-102 and we may dismiss it because of the large impact parameter. The other example, however, is for a $z = 0.0081$ early-type galaxy that has the lowest impact parameter ($\rho = 34 \text{ kpc}$) of the entire sample! Furthermore, it is the only example we have where the sightline is certain to intersect the galaxy's virialized halo. We identify no other galaxies at this redshift and only note that it might be associated with the extreme outskirts of Virgo. Either way, this non-detection stresses that the IGM/galaxy connection shows special cases that contradict dominant trends.

The lower panels of Figure 1 present the equivalent width of O VI 1031 (W^{1031} ; when UV spectral coverage exists) and an estimate of the O⁺ column density N_{OVI} for the strongest absorber within 300 km s⁻¹ of each galaxy. In contrast to Ly α , the results are dominated by non-detections. Furthermore, 3 of 5 systems with a positive detection (3/4 for those with $N_{\text{OVI}} > 10^{14} \text{ cm}^{-2}$) also show a brighter galaxy ($L > 0.1L^*$) within 150 kpc of the sightline. Ignoring these systems, 0/13 of 'isolated' dwarf galaxies show a positive detection to an equivalent width limit of 200 mÅ and only 2/8 for a 50 mÅ limit. We conclude that the CGM of dwarf galaxies at impact parameters of 50–300 kpc rarely shows O VI absorption characteristic of current UV surveys. The CGM of dwarf galaxies does give rise to nearly ubiquitous H I absorption, but has insufficient surface density, metallicity, and/or the physical conditions (i.e. density, temperature) needed to also exhibit significant O VI absorption.

In summary, the circumgalactic medium of dwarf galaxies at $\rho \gtrsim 100 \text{ kpc}$ is characterized by a high incidence of moderate strength Ly α absorption ($W^{\text{Ly}\alpha} \gtrsim$




FIG. 2.— Same as Figure 1 but for L^* galaxies. Note the high covering fraction to $\text{Ly}\alpha$ absorption (with column densities typically exceeding 10^{14} cm^{-2}) but the lack of O VI detections for $\rho > 225 \text{ kpc}$ to sensitive limits. The results suggest that O VI gas near L^* galaxies is primarily associated with their virialized halos but not their extended CGM environments.

200 mÅ) and a low incidence of O VI absorption to modest equivalent width limits. Our survey, unfortunately, offers few probes of dwarf galaxies on smaller scales, i.e. with sightlines intersecting their dark matter halos. This assessment awaits a dedicated survey of such galaxies.⁷ **L^* Galaxies:** In Figure 2, we plot the same quantities versus impact parameter but now for $L > L^*$ galaxies (Table 4). Our survey has only a modest sample of such bright galaxies within 300 kpc of the sightlines which reflects, of course, their lower comoving number density. Similar to the dwarf galaxies, nearly every L^* galaxy exhibits significant $\text{Ly}\alpha$ absorption.⁸ This even includes luminous, early-type galaxies which are likely gas-poor on galactic-scales. The average equivalent width at $\rho = 100\text{--}200 \text{ kpc}$ is $W^{\text{Ly}\alpha} \approx 500 \text{ mÅ}$ and $\approx 400 \text{ mÅ}$ at $\rho = 200\text{--}300 \text{ kpc}$, twice that observed for the dwarf galaxies (ignoring dwarfs with bright neighbors; Table 3). Similarly, the N_{HI} values of the L^* galaxies exceed those for the dwarfs. This implies that

the H I gas near a typical L^* galaxy has a higher total surface density, neutral fraction, and/or a more extreme velocity field. Such trends with luminosity have been revealed previously (e.g. Chen et al. 2001b); presumably, they are a simple consequence of these galaxies having greater mass.

Regarding O VI absorption, the L^* galaxies show a greater fraction (5/10) of positive detections than dwarf galaxies. Furthermore, there is a possible trend with impact parameter: all of the detections occur within $\rho = 225 \text{ kpc}$, which corresponds (roughly) to the virial radii of these luminous galaxies. The figure also reveals that every emission-line galaxy exhibits a positive O VI detection; all of the non-detections are associated with early-type spectra. A preference for O VI absorption to occur in late-type galaxies was previously reported by Chen & Mulchaey (2009) for their modest sample of galaxy/O VI associations (Tumlinson et al., in prep.). Unfortunately, all of our non-detections also occur at large ρ and, therefore, we cannot separate the effects of impact parameter and spectral type with this sample. Nevertheless, the results demonstrate clearly that the extended CGM of L^* galaxies (at least those with early-type spectra) is frequently characterized by the non-detection of O VI gas.

⁷ An ongoing Cycle 18 HST/COS survey by J. Tumlinson (GO 12248) is designed to address this directly, although it will not include high S/N spectra of the O VI doublet.

⁸ The only notable exception is an $L \approx 1.0 L^*$ galaxy with an early-type spectrum located at an impact parameter of $\rho \approx 250 \text{ kpc}$ from PKS0405–123 ($z = 0.203$). Perhaps not coincidentally, it also is associated with one of the most significant galaxy overdensities identified in our survey.

TABLE 4
 $L > L^*$ GALAXIES WITHIN 300 KPC OF A QSO SIGHTLINE

Field	RA (J2000)	DEC (J2000)	z_{gal}	T^a	ρ (kpc)	L (L^*)	$\mathcal{N}^b$	$z_{\text{abs}}^{\text{Ly}\alpha}$	$W^{\text{Ly}\alpha}$ (mÅ)	N_{HI}	Rf.	$z_{\text{abs}}^{\text{OVI}}$	W^{OVI} (mÅ)	N_{OVI}	Rf.
Q0026+1259	00:29:15.36	+13:20:57.0	0.0393	E	213	1.384	0	...	...	...	...	0.0393	< 120	< 14.0	9
PKS0405-123	04:07:51.28	-12:11:38.3	0.1670	L	115	2.143	1	0.1671	697	15.47	3	0.1669	412	14.64	1
	04:07:42.79	-12:11:33.1	0.2030	E	262	1.070	9	0.2030	< 50	< 13.00	9	0.2030	< 30	< 13.4	9
	04:07:50.69	-12:12:25.2	0.2976	E	245	1.878	7	0.2977	264	13.85	3	0.2976	< 50	< 13.6	9
	04:07:45.96	-12:11:09.9	0.3612	E	213	6.604	5	0.3608	554	15.14	3	0.3616	96	14.00	1
PG1004+130	10:07:34.55	+12:52:09.5	0.0704	E	290	2.404	0	...	...	...	...	0.0704	< 100	< 13.9	9
PG1116+215	11:19:06.73	+21:18:29.3	0.1383	E	129	2.039	4	0.1385	479	> 14.35	1	0.1385	83	13.95	1
PKS1302-102	13:05:20.22	-10:36:30.4	0.0426	L	212	2.577	1	0.0422	410	14.83	3	0.0422	194	14.38	3
FJ2155-0922	21:54:56.64	-09:18:07.9	0.0517	E	251	1.427	0	0.0515	272	14.08	3	0.0514	< 47	< 13.6	4
	21:55:06.53	-09:23:25.2	0.1326	L	212	1.626	2	0.1324	487	> 14.30	1	0.1324	225	14.39	1

REFERENCES. — 1: Tripp et al. (2008); 2: Thom & Chen (2008b); 3: Danforth & Shull (2008); 4: Danforth et al. (2006); 5: Penton et al. (2004); 6: Chen & Mulchaey (2009); 9: This paper.

^a Spectral type (E=Early-type; L=Late-type; see text for the quantitative definition).

^b Number of additional galaxies within 3 Mpc of the sightline, 400 km s⁻¹ of this galaxy, and having $L > 0.1L^*$.




FIG. 3.— Same as for Figure 1 but for sub- L^* galaxies. One notes a strong correlation between $W^{\text{Ly}\alpha}$ and impact parameter for these galaxies and also an apparent division at $\rho \approx 100$ kpc between $W^{\text{Ly}\alpha} \gtrsim 1 \text{ \AA}$ and lines exhibiting a large scatter of lower values. Intriguingly, this division occurs near the expected virial radius for these galaxies. In contrast to the dwarf and L^* galaxies (Figures 1–2), we observe a very high detection rate for O VI at all impact parameters. This indicates the extended CGM of sub- L^* galaxies has a high covering fraction to modest O VI absorption. The gray diamond indicates the one sub- L^* galaxy in our sample that has an L^* neighbor within 150 kpc of the sightline.

Sub- L^* Galaxies: Our final galaxy subset is the sub- L^* galaxies ($0.1L^* < L < L^*$; Table 5). Figure 3 reveals a 100% detection rate for H I gas to $\rho = 300$ kpc for these galaxies. In contrast to the other galaxy subsets, the Ly α equivalent widths of the sub- L^* galaxies show a very obvious trend with impact parameter. A Spearman’s test reports that the null hypothesis of no correlation is ruled out at $> 99.5\%$ c.l. There is also the hint of a division in the $W^{\text{Ly}\alpha}$ values at $\rho \approx 100$ kpc, i.e. between exclusively large equivalent widths ($W^{\text{Ly}\alpha} > 1 \text{ \AA}$ for $\rho < 100$ kpc) and a large scatter of primarily lower values for $\rho > 100$ kpc (Table 6). As this impact parameter roughly coincides with the virial radius expected for sub- L^* galaxies, if confirmed, it may be related to galactic-scale processes at $\rho < r_{\text{vir}}$ whereas the $W^{\text{Ly}\alpha}$ values at larger ρ trace the physical properties of the surrounding (unvirialized) CGM.

Because the Ly α line is saturated for most of the systems, the $W^{\text{Ly}\alpha}$ values are a good proxy for kinematics, i.e. the systems at $\rho < 100$ kpc require a velocity spread of $\Delta v = cW^{\text{Ly}\alpha}/(1215.67 \text{ \AA}) \approx 250 \text{ km s}^{-1} (W^{\text{Ly}\alpha}/1 \text{ \AA})$. Such motions may be just consistent with galaxies having circular velocities $v_c \gtrsim 100 \text{ km s}^{-1}$, a reasonable estimate for sub- L^* galaxies. There is also the hint of a difference in Ly α equivalent width for galaxies with differing spectral type: the late-type galaxies have systematically higher $W^{\text{Ly}\alpha}$ values than early-type galaxies at a similar impact parameter. This conclusion is tempered, however, by the relatively small sample size and should be confirmed with a larger dataset.

Remarkably, the sub- L^* galaxies show a very high incidence of associated O VI absorption for impact parameters $\rho < 300$ kpc. At $\rho < 100$ kpc, 3/4 (75%) of the galaxies have $W^{1031} > 50 \text{ m\AA}$ ($N_{\text{OVI}} > 10^{13.5} \text{ cm}^{-2}$). And at larger impact parameters, the incidence is yet higher. It appears the association extends well beyond the virial radii of these galaxies; the results require that sub- L^* galaxies are surrounded by a CGM that bears O VI gas to an impact parameter of several hundreds kpc. We also emphasize that only one of the galaxies has a neighboring, bright ($L > L^*$) galaxy close to the sightline. Finally, we note a possible trend of decreasing W^{1031} (and N_{OVI}) with increasing impact parameter. This trend, however, is dominated by the two cases at $\rho \approx 100$ kpc with large W^{1031} values.

The non-detection of O VI absorption for the galaxy at $z = 0.081$ and $\rho = 33$ kpc in the FJ2155–0922 field warrants additional discussion. It is striking, given the high incidence of O VI detections, that the galaxy at smallest impact parameter would exhibit the lowest W^{1031} value.⁹ Furthermore, as is evident from Figure 3 and Table 5, there is very strong Ly α absorption associated with this galaxy ($W^{\text{Ly}\alpha} > 1 \text{ \AA}$). It has an early-type spectrum, and its luminosity places it at the upper end of the sub- L^* population. It is noteworthy that this galaxy lies within an overdensity in the FJ2155–0922 field: 10 galaxies with $L > 0.1L^*$ lie within 3 Mpc of the sightline. On the other hand, the $z = 0.0788$, sub- L^* galaxy ($\rho = 221$ kpc) in the same field has 22 such neighbors yet shows modest O VI absorption. Similar to the non-detection of Ly α at

$\rho \approx 30$ kpc for the dwarf galaxy in the PG1216+069 field, this one non-detection of O VI gas stresses the complexity of galaxy/absorber associations.

In summary, all three classes of galaxies considered above exhibit high covering fractions to Ly α absorption for impact parameters $\rho < 300$ kpc. There are possible trends of increasing $W^{\text{Ly}\alpha}$ values for brighter galaxies and decreasing $W^{\text{Ly}\alpha}$ values at higher impact parameter. In contrast, the covering fraction to O VI absorption is highest for sub- L^* galaxies, low for dwarf galaxies, and low for early-type (and/or high- ρ) L^* galaxies. These results are empirical descriptions of the gas associated with low- z galaxies; we explore further the implications for the origins of Ly α and O VI absorption in the following sections.

3.2. Galaxies Associated with IGM Absorption

In the previous sub-section, we studied the Ly α and O VI absorption associated with galaxies at low impact parameters to quasar sightlines. For that analysis, the measures (luminosity, impact parameter, equivalent width) were all well-defined and the results were relatively insensitive to biases related to incompleteness in the galaxy survey. The key complication of such analysis, however, is that galaxies cluster and therefore their environment (e.g. the presence of a brighter, nearby neighbor) may play a significant role in observed associations between a galaxy and the IGM. Furthermore, this approach only indirectly addresses fundamental questions related to the origin of Ly α and O VI gas in the IGM.

One may gain further insight by reversing the experiment, i.e. to study the properties of the closest (detected) galaxy to a given set of IGM absorption lines. In this manner, one tests how frequently galaxies and their halos are associated with various components of the IGM, and one may assess their physical properties and environment. This approach, however, has the obvious downside that no galaxy survey is truly complete, neither in terms of field-of-view nor depth. Indeed, our survey does not cover sufficient area to identify all galaxies within $\rho \approx 200$ kpc for $z < 0.02$ and our magnitude limit implies significant incompleteness to $L < 0.1L^*$ in most fields for $z > 0.1$. Therefore, we restrict the following analysis and discussion to the redshift interval $0.02 < z < 0.2$ and caution that at the high end we have limited sensitivity to very faint galaxies.

In Figure 4, we present the analysis for Ly α where the symbols label the luminosity of the closest galaxy detected and the upward arrows¹⁰ indicate absorbers without an associated galaxy in our survey. For very low column densities, $10^{13} \text{ cm}^{-2} < N_{\text{HI}} < 10^{14} \text{ cm}^{-2}$, approximately 40% of the absorbers may be associated with a galaxy within 1 Mpc, always at $\rho > 100$ kpc and primarily at $\rho > 200$ kpc (80% of the associations). We conclude that *few, if any, of the weak Ly α absorbers arise from the virialized halos of $z \sim 0$ galaxies*. Instead, weak H I absorption traces either the extended CGM of galaxies or, more commonly, structures with a low filling factor of galaxies (non-detections). The results are similar for the Ly α lines with modest N_{HI} (10^{14} – 10^{15} cm^{-2}), al-

⁹ We have confirmed the measurements of Danforth et al. (2006) by analyzing our extraction of the *FUSE* spectrum for FJ2155–0922.

¹⁰ These arrows are placed at the physical distance equivalent to $10'$ angular separation (the typical field-of-view for the LCO/WFCCD survey) at the redshift of the absorber.

TABLE 5
SUB- L^* GALAXIES ($0.1L^* < L < L^*$) WITHIN 300 KPC OF A QSO SIGHTLINE

Field	RA (J2000)	DEC (J2000)	z_{gal}	T^a	ρ (kpc)	L (L^*)	$\mathcal{N}^b$	$z_{\text{abs}}^{\text{Ly}\alpha}$	$W^{\text{Ly}\alpha}$ (mÅ)	N_{HI}	Rf.	$z_{\text{abs}}^{\text{OVI}}$	W^{OVI} (mÅ)	N_{OVI}	Rf
TONS180	00:57:08.52	-22:18:29.6	0.0456	L	265	0.365	1	0.0456	212	13.80	5	0.0456	62	13.74	4
PKS0312-77	03:12:01.76	-76:55:17.7	0.0594	L	245	0.292	0	0.0595	153	13.53	3	...	...	...	...
PKS0405-123	03:11:57.89	-76:51:55.6	0.2026	L	126	0.634	2	0.2027	< 952	15.14	3	0.2027	538	> 14.9	1
	04:07:58.12	-12:12:24.7	0.0965	L	255	0.150	8	0.0966	484	14.64	3	0.0966	59	13.71	3
	04:07:44.03	-12:12:09.5	0.1532	E	181	0.980	1	0.1522	128	13.46	3	0.1522	< 50	< 13.6	9
PG1116+215	04:07:48.40	-12:12:10.8	0.3520	U	156	0.273	2	0.3509	380	14.04	3	0.3509	24	13.35	3
	11:19:05.55	+21:17:33.3	0.0600	E	124	0.104	5	0.0593	187	13.64	1	0.0593	63	13.77	1
	11:19:12.20	+21:18:52.0	0.1660	E	155	0.336	15	0.1655	765	> 14.4	1	0.1655	111	14.08	1
PG1211+143	12:14:09.54	+14:04:21.3	0.0511	L	130	0.829	8	0.0510	703	15.67	3	0.0513	...	14.30	4
	12:14:19.86	+14:05:10.3	0.0644	E	140	0.740	0	0.0645	535	15.73	3	0.0645	144	14.16	3
	12:19:23.45	+06:38:20.3	0.1241	L	87	0.650	6	0.1236	1433	> 14.8	1	0.1236	378	14.70	1
PKS1302-102	13:05:25.65	-10:39:23.5	0.0420	L	294	0.328	1	0.0422	410	14.83	3	0.0422	194	14.38	3
	13:05:32.19	-10:33:56.9	0.0936	E	64	0.195	7	0.0949	681	15.35	3	0.0948	84	13.75	4
	13:05:35.30	-10:33:24.2	0.1453	L	83	0.396	2	0.1453	890	15.29	3	0.1453	146	14.16	4
FJ2155-0922	13:05:34.97	-10:34:22.6	0.1917	E	209	0.888	2	0.1916	401	15.01	3	0.1916	80	13.93	1
	21:54:50.87	-09:22:33.3	0.0788	L	221	0.157	22	0.0777	448	> 14.3	1	0.0777	46	13.61	1
	21:54:59.96	-09:22:24.7	0.0810	E	32	0.872	10	0.0808	1007	15.11	3	0.0807	< 14	< 13.1	4
	21:55:01.62	-09:20:47.0	0.1555	E	246	0.441	1	0.1549	63	13.06	3	...	...	...	...

REFERENCES. — 1: Tripp et al. (2008); 2: Thom & Chen (2008b); 3: Danforth & Shull (2008); 4: Danforth et al. (2006); 5: Penton et al. (2004); 6: Chen & Mulchaey (2009); 9: This paper.

^a Spectral type (E=Early-type, L=Late-type, U=unknown; see text for the quantitative definition).

^b Number of additional galaxies within 3 Mpc of the sightline, 400 km s⁻¹ of this galaxy, and having $L > 0.1L^*$.

TABLE 6
GALAXY/ABSORBER STATISTICS

Luminosity	ρ (kpc)	$\mathcal{N}$	Med($W^{\text{Ly}\alpha}$) (Å)	RMS($W^{\text{Ly}\alpha}$) (Å)	$\mathcal{N}$	M(N_{HI})	R(N_{HI})	$\mathcal{N}$	M(W^{1031}) (mÅ)	R(W^{1031}) (mÅ)	$\mathcal{N}$	M(N_{OVI})	R(N_{OVI})
$L < 0.1L^*$	0-100	7	0.37	0.77	7	15.4	2.0	8	<200	121	8	<14.2	0.4
	100-200	4	0.22	0.29	4	13.8	1.1	4	< 43	84	5	<13.5	0.6
	200-300	8	0.28	0.28	8	14.0	0.5	7	< 45	36	7	<13.6	0.4
$0.1 < L < L^*$	0-100	4	1.01	0.32	4	15.3	0.3	4	146	158	4	14.2	0.7
	100-200	7	0.54	0.31	7	14.4	0.9	6	111	192	7	14.1	0.5
	200-300	7	0.40	0.16	7	14.3	0.7	5	62	60	5	13.7	0.3
$L > L^*$	0-100	2	0.70	0.15	2	15.5	0.8	2	412	232	2	14.6	0.5
	100-200	6	0.41	0.18	6	14.3	0.8	8	<100	70	8	<14.0	0.4
	200-300	6	0.41	0.18	6	14.3	0.8	8	<100	70	8	<14.0	0.4
All Galaxies	0-100	11	0.68	0.64	11	15.3	1.6	12	<146	127	12	<14.2	0.5
	100-300	26	0.28	0.23	26	14.0	0.7	25	< 62	107	26	<13.8	0.4
	300-500	32	0.16	0.14	31	13.7	0.7	18	< 44	33	21	<13.6	0.4
	500-1000	39	0.05	0.16	38	13.0	0.8	17	< 40	36	18	<13.6	0.4

NOTE. — For column densities, all statistics are calculated on log N . For the columns, $\mathcal{N}$ gives the number of galaxies, M() refers to median statistic, and R() refers to RMS.

^a Spectral type (see text for the quantitative definition).

^b Impact parameter and luminosity of the brightest galaxy ($L > 0.1L^*$) within 200 kpc and $|\delta v| < 300 \text{ km s}^{-1}$ (if any).

though the incidence of an association to a galaxy with $\rho < 1 \text{ Mpc}$ is higher (70%). For the small set of strong H I absorption lines ($N_{\text{HI}} > 10^{15} \text{ cm}^{-2}$), the results are qualitatively different. Every Ly α line is linked to a galaxy and the majority of these have $\rho < 200 \text{ kpc}$. Even in these cases, however, the association with a virialized halo may be tenuous. Altogether, our results do offer further evidence that large N_{HI} absorbers are physically associated with galaxies whereas weak Ly α lines have no direct physical connection (Chen et al. 2005; Shone et al. 2010).

Turning to O VI absorption, Figure 5 shows the galaxy at the closest impact parameter to the 30 published O VI

absorbers along our quasar sightlines with $0.02 < z < 0.2$ (Table 7). This includes 5 O VI systems where we have not identified a galaxy within 1 Mpc of the sightline with $|\delta v| < 400 \text{ km s}^{-1}$. The rate of non-detections is much lower¹¹ than that observed for weak Ly α lines. Of course, only a few of the O VI systems have low H I column densities ($N_{\text{HI}} < 10^{14} \text{ cm}^{-2}$). Ignoring the non-detections, the median impact parameter to a galaxy is approximately 200 kpc (consistent with the results of

¹¹ This lends further support to the assertion that the lower fraction of weak H I absorbers associated with galaxies is not driven by incompleteness in the LCO/WFCCD galaxy survey.




FIG. 4.— Impact parameter (symbols indicate luminosity) of the galaxy closest to the sightline with a redshift coincident ($|\delta v| < 400 \text{ km s}^{-1}$) to the Ly α absorbers detected along our quasar sightlines. Note that most of the N_{HI} values exceeding $10^{14.5} \text{ cm}^{-2}$ should be considered lower limits due to line saturation. The dotted lines (from bottom to top) mark estimates for the virial radii of the dwarf, sub- L^* , and L^* galaxies, respectively. In cases where we have not detected a galaxy within 1 Mpc of the sightline, we plot an upward arrow at the distance at each absorber’s redshift corresponding to the typical $10'$ field-of-view for our LCO/WFCCD galaxy survey. For $N_{\text{HI}} < 10^{14.5} \text{ cm}^{-2}$, the incidence of associated galaxies is low and those that are detected generally lie at $\rho > 200 \text{ kpc}$. We conclude that few, if any, of these absorbers arise in the virialized halos of $z \sim 0$ galaxies. At higher column densities, the fraction of Ly α absorbers with an associated galaxy increases and the impact parameter to the systems decreases suggesting a physical association between the gas and galaxy.

Stocke et al. 2006). This offset is smaller than that observed for Ly α lines having $N_{\text{HI}} \approx 10^{14} \text{ cm}^{-2}$. On the other hand, only a small fraction of the O VI systems (5/30) have a galaxy detected within 100 kpc and only 1/18 for $N_{\text{OVI}} < 10^{14} \text{ cm}^{-2}$. There are very few cases in the sample where one would insist that the sightline has intersected the virialized halo of the associated galaxy. Instead, the O VI gas appears linked to the extended CGM of galaxies on scales of a few hundred kpc. The luminosities of the nearest neighbors include galaxies at a wide range of luminosity and spectral type, with a preference toward late-type sub- L^* systems. There is an apparent trend of decreasing ρ with increasing N_{OVI} for the range of values considered in our analysis, but this is tempered by the large scatter in these quantities.

As emphasized at the start of this sub-section, the results presented in Figures 4 and 5 are subject to the incompleteness of our galaxy survey. In particular, one may argue that a population of fainter galaxies lie at smaller impact parameters than the sample comprising our survey. This could significantly affect our infer-

ences regarding the roles of virialized halos and the extended CGM in hosting Ly α /O VI absorption. To assess this concern, Table 7 lists the luminosity limit for each absorber and the completeness of the galaxy survey to that limit for $\rho = 150 \text{ kpc}$. Regarding completeness, with the exception of two fields (PG1116+215 and PG1211+143), we obtained a redshift for every galaxy within $\rho = 150 \text{ kpc}$ of the O VI absorber to the magnitude limit $R = 19.5 \text{ mag}$. We are confident, therefore, that no galaxy of comparable luminosity lies at closer distance to the sightline than the ones we have discovered.¹² Regarding the luminosity limit, a subset of the galaxies lie within twice the limit yet $\approx 40\%$ are $5\times$ brighter than the formal limit. Furthermore, owing to the details of mask design and specific scientific interests, we have surveyed several fields to $R > 19.5 \text{ mag}$

¹² Although we are not sensitive to galaxies at very small impact parameters to the quasar (i.e. $\theta \lesssim 1''$, or $\rho \lesssim 20 \text{ kpc}$ for the redshifts considered), we expect these to be very rare occurrences and note that none of the sightlines exhibit very large H I column densities ($N_{\text{HI}} \gg 10^{17} \text{ cm}^{-2}$).




FIG. 5.— Impact parameter (symbols indicate luminosity) of the galaxy closest to the sightline with a redshift coincident ($|\delta v| < 400 \text{ km s}^{-1}$) to the O VI absorbers detected along our quasar sightlines. The dotted horizontal lines mark estimates (from bottom to top) for the virial radii of the dwarf, sub- L^* , and L^* galaxies, respectively. The arrows are positioned as in Figure 4. One notes that although most of the O VI absorbers have an associated galaxy within a few hundred kpc of the sightline, very few of the sightlines intersect the galaxy’s expected virialized halo. The results suggest that O VI gas is associated primarily with the extended CGM of $z \sim 0$ galaxies. There is possibly an anti-correlation between ρ and N_{OVI} , but the significance of such a trend is tempered by the large scatter.

albeit with lower completeness. The galaxies discovered to these fainter magnitudes have been included in the analysis. We conclude, therefore, that our results are unlikely to be severely compromised by a lack of sensitivity to fainter galaxies.

Another (brute force) approach to assess the effects of incompleteness is to obtain significantly deeper spectroscopy in the fields. Chen & Mulchaey (2009) have performed a galaxy survey with the IMACS and LDSS3 spectrometers on the 6.5 m Magellan telescopes for two of our fields (PKS0405–123, PG1216+069). They achieved a high completeness around each quasar sightline to several magnitudes fainter limits ($\approx 10\times$ deeper). For the PKS0405–123 field, they do identify a new dwarf galaxy associated with the $z = 0.0918$ O VI absorber which lies at closer impact parameter ($\rho = 73 \text{ kpc}$) than the dwarf galaxy we had discussed. This is, however, the only modification for the four O VI absorbers within the redshift interval $0.02 < z < 0.2$, and even this galaxy is at an impact parameter that may exceed its virialized halo. For the PG1216+069 field, Chen & Mulchaey (2009) find no fainter galaxies at closer impact parameters for the $z = 0.125$ sub- L^* galaxy identified in our survey.

4. DISCUSSION

The previous section described the association of Ly α and O VI absorption with galaxies discovered in our LCO/WFCCD survey (Paper IV). In § 3.1, we explored the strength of Ly α and O VI absorption as a function of galaxy luminosity, spectral type, and impact parameter. We then reported on the results of a search for galaxies related to Ly α and O VI absorption (§ 3.2). In this section, we synthesize these results and discuss the implications in the context of previous observational and theoretical work. We divide the discussion by transition (i.e. Ly α and O VI) but emphasize that a complete description of the IGM must consider both elements.

4.1. Ly α

The Ly α transition is the strongest and most commonly observed transition of the intergalactic medium. Its complex pattern of absorption defines the so-called Ly α forest and sets the starting point for nearly all IGM analysis. The current cosmological paradigm for the Ly α forest is that these lines arise from an undulating Gunn-Peterson field of overdensities (e.g. Miralda-Escudé et al. 1996). In turn, observations of the IGM provide direct constraints for cosmological param-

TABLE 7
O VI ABSORBERS IDENTIFIED IN OUR FIELDS

Field	z_{OVI}	W^{OVI} (mÅ)	N_{OVI}	Ref	z_n^a	ρ_n^a (kpc)	L_n^a (L^*)	L_{lim}^b (L^*)	C^c
0.005 < z < 0.02									
PG1116+215	0.0050	...	13.91	4	...	...	...	0.000	66
PG1211+143	0.0071	...	13.78	3	...	...	...	0.000	52
3C273	0.0033	31	13.41	1	...	...	...	0.000	73
3C273	0.0076	14	13.11	3	...	...	...	0.001	74
FJ2155-0922	0.0173	< 350	13.90	4	...	...	...	0.003	100
0.02 < z < 0.2									
TONS180	0.0456	62	13.74	4	0.0456	265	0.377	0.020	100
TONS180	0.0234	43	13.48	4	0.0234	148	0.019	0.005	100
TONS180	0.0436	31	13.43	4	...	...	...	0.018	100
PKS0312-77	0.1983	64	13.84	1	0.1982	680	0.785	0.464	100
PKS0312-77	0.1589	86	13.94	1	...	...	...	0.284	100
PKS0405-123	0.0918	49	13.58	4	0.0906	410	0.067	0.087	100
PKS0405-123	0.0966	59	13.71	3	0.0965	256	0.150	0.097	100
PKS0405-123	0.1669	412	14.64	1	0.1670	93	0.080	0.316	100
PKS0405-123	0.1829	84	13.94	1	...	...	...	0.387	100
PG1116+215	0.1655	111	14.08	1	0.1660	156	0.336	0.311	80
PG1116+215	0.1385	83	13.95	1	0.1383	130	2.039	0.210	80
PG1116+215	0.0593	63	13.77	1	0.0600	124	0.104	0.035	67
PG1211+143	0.0513	...	14.30	4	0.0511	131	0.829	0.026	64
PG1211+143	0.0645	144	14.16	3	0.0646	69	0.084	0.041	75
PG1216+069	0.1236	378	14.70	1	0.1241	88	0.650	0.164	100
3C273	0.0902	16	13.13	1	0.0902	443	1.105	0.084	100
3C273	0.1200	24	13.33	1	...	...	...	0.154	100
PKS1302-102	0.0940	19	13.15	4	0.0936	65	0.195	0.091	100
PKS1302-102	0.0647	69	13.70	3	0.0647	338	0.049	0.042	100
PKS1302-102	0.0989	...	14.03	3	...	...	...	0.102	100
PKS1302-102	0.0423	194	14.38	3	0.0426	212	2.577	0.017	80
PKS1302-102	0.1453	146	14.16	4	0.1453	83	0.396	0.233	100
PKS1302-102	0.1916	80	13.93	1	0.1917	209	0.888	0.430	100
MRK1383	0.0519	66	13.83	4	...	...	...	0.026	60
FJ2155-0922	0.1579	120	14.08	1	0.1581	336	0.495	0.280	100
FJ2155-0922	0.1324	225	14.39	1	0.1326	212	1.626	0.191	100
FJ2155-0922	0.0776	46	13.61	1	0.0788	222	0.157	0.061	100
FJ2155-0922	0.1765	...	14.44	1	0.1764	331	2.399	0.358	100
PKS2155-304	0.0571	44	13.57	3	0.0570	406	1.595	0.032	100
PKS2155-304	0.0541	32	13.63	3	0.0541	544	2.053	0.029	100
$z > 0.2$									
PKS0312-77	0.2027	538	14.91	1	0.2026	127	0.634	0.487	100
PKS0405-123	0.4950	213	14.44	1	...	...	...	3.855	100
PKS0405-123	0.3509	24	13.35	3	0.3520	157	0.273	1.707	100
PKS0405-123	0.3616	96	14.00	1	0.3612	214	6.604	1.831	100
PKS0405-123	0.3634	38	13.61	1	...	...	...	1.852	100
PG1216+069	0.2677	22	13.30	1	...	...	...	0.913	100
PG1216+069	0.2819	94	13.95	1	...	...	...	1.028	100
PKS1302-102	0.2274	40	13.59	1	...	...	...	0.631	100
PKS1302-102	0.2256	82	13.93	1	0.2256	417	0.892	0.620	100
PKS1302-102	0.2044	33	13.64	3	...	...	...	0.496	100
PKS1302-102	0.2529	11	12.92	3	...	...	...	0.802	100

NOTE. — The analysis presented in § 3.2 only considers O VI systems with $0.02 < z < 0.2$ to minimize incompleteness in the galaxy survey.

^a Impact parameter and luminosity of the brightest galaxy ($L > 0.1L^*$) within 200 kpc and $|\delta v| < 300 \text{ km s}^{-1}$ (if any).

^b Impact parameter and luminosity of the brightest galaxy ($L > 0.1L^*$) within 200 kpc and $|\delta v| < 300 \text{ km s}^{-1}$ (if any).

^c Completeness of the galaxy survey to $R = 19.5 \text{ mag}$ within the radius corresponding to $\rho = 150 \text{ kpc}$ at the O VI absorber's redshift.

ters (e.g. Dunkley et al. 2009). This gas is also predicted to be the dominant baryonic reservoir of the universe, which fuels the formation and growth of galaxies.

We begin with a discussion of the properties of H I gas surrounding low- z galaxies ($\rho < 300$ kpc). Several previous studies have examined the incidence of Ly α absorption for galaxies located at small impact parameters to quasar sightlines. In each case, the authors have reported a very high detection rate for galaxies within a few hundred kpc: Chen et al. (2001a) found positive detections for 29/31 of their predominantly L^* galaxies to $\rho = 250 h_{72}^{-1}$ kpc with $W^{\text{Ly}\alpha} > 300$ mÅ and reported a sharp decline at larger impact parameter; Bowen et al. (2002) detected Ly α lines with $N_{\text{HI}} \geq 10^{13} \text{ cm}^{-2}$ for 8/8 of the galaxies located within $\rho = 280 h_{72}^{-1}$ kpc of their quasar sightlines, and Wakker & Savage (2009) reported positive detections for 7/7 of the $L \geq 0.1 L^*$ field galaxies in their sample when restricting to $\rho \leq 350$ kpc.

With our LCO/WFCCD survey we have examined a sample of 37 galaxies at $\rho < 300$ kpc with greater emphasis on lower luminosities than previous work. The results presented in Figure 1–3 (Tables 3–5) further solidify the correlation between galaxies and Ly α absorption: to an impact parameter of $\rho \approx 300$ kpc, we observe a nearly 100% probability for detecting Ly α absorption ($W^{\text{Ly}\alpha} > 50$ mÅ) within 400 km s^{-1} of a galaxy. This result is independent of galaxy luminosity (for $L > 0.01 L^*$), spectral type, and local environment¹³ (defined by the number of close neighbors). Altogether, these results demand that low- z galaxies of all types are embedded within a diffuse, highly ionized medium with a very high covering fraction to significant H I absorption. We refer to this medium as the circumgalactic medium (CGM), although this gas need not be physically associated (e.g. bound) to the galaxy.

Before considering the origin of this medium, we review and then explore its physical properties. First, we note that the Doppler parameters that are typical of low- z Ly α lines generally require kinetic temperatures $T < 10^5$ K (e.g. Davé et al. 1999; Williger et al. 2006). The standard expectation is that the majority of these absorbers trace photoionized gas with $T \sim 10^4$ K. For those absorption systems that also exhibit metal-line transitions (e.g. C IV, C III, O VI), this hypothesis may be tested against ionization modeling and the results are usually consistent with a photoionized medium (e.g. Prochaska et al. 2004; Cooksey et al. 2008). This modeling also provides a constraint on the ionization fraction and an estimate of the total hydrogen column density N_{H} .

A remarkable result for the low- z IGM is that the estimated N_{H} values are uniform at $\approx 10^{19} \text{ cm}^{-2}$, i.e., independent of the observed N_{HI} value (Prochaska et al. 2004; Lehner et al. 2007). This ‘ N_{H} constancy’ was interpreted by Prochaska et al. (2004) to result from lower N_{HI} absorbers having lower average volume density (e.g. Davé et al. 1999) which implies a higher ionization correction for a uniform (background) radiation field. Indeed, analysis of recent simulations of the low- z IGM

have considered the N_{H} constancy explicitly and have reproduced the observed result with a similar explanation (Davé et al. 2010). We caution that this N_{H} constancy was derived empirically from absorbers with $N_{\text{HI}} \approx 10^{13.5} - 10^{15} \text{ cm}^{-2}$ which had no known association to galaxies. We demonstrate below, however, that the majority of the stronger N_{HI} systems must be associated with galaxies.

Adopting a constant N_{H} value for the observed Ly α forest, we may crudely estimate the mass of the photoionized CGM surrounding low- z galaxies. A simple estimate follows from the characteristic hydrogen column density $N_{\text{H,CGM}}$ and a characteristic radius r_{CGM} :

$$M_{\text{CGM}} = N_{\text{H,CGM}} m_p \mu \pi r_{\text{CGM}}^2, \quad (2)$$

with the factor $\mu \approx 1.3$ accounting for helium. For our assumed values, we find

$$M_{\text{CGM}} \approx 3 \times 10^{10} M_{\odot} \left(\frac{N_{\text{H,CGM}}}{10^{19} \text{ cm}^{-2}} \right) \left(\frac{r_{\text{CGM}}}{300 \text{ kpc}} \right)^2. \quad (3)$$

Therefore, we estimate a baryonic reservoir surrounding each galaxy which meets or significantly exceeds the typical baryonic mass of present-day sub- L^* galaxies. Again, this CGM need not be physically associated (e.g. gravitationally bound) with the galaxy. Nevertheless, it represents a large baryonic reservoir for future star formation or could even (in part) be gas that was expelled during the processes of galaxy formation. Lastly, if the gas is metal enriched, the mass in metals could exceed the amount metals locked in stars within these galaxies.

The near unity covering fraction of this CGM for Ly α absorption to $\rho \approx 300$ kpc inspires us to examine further its radial extent. Previously, Chen et al. (2001b) reported a decline in the average Ly α equivalent width for $\rho \gtrsim 250$ kpc. Similarly, Tripp et al. (1998) and Wakker & Savage (2009) noted a lower detection rate and average Ly α equivalent width for $L \approx L^*$ galaxies to $\rho \gtrsim 1$ Mpc. In Figure 6 (Table 8), we have extended the analysis of our LCO/WFCCD survey to $\rho = 1$ Mpc. Here we include galaxies of all luminosity and plot only the closest galaxy to the sightline without repeating any pair of galaxies with $|\delta v| < 400 \text{ km s}^{-1}$. We also include the galaxy survey in the field of PKS0405–123 performed by Williger et al. (2006). There are two issues to emphasize before proceeding: (i) incompleteness in our galaxy survey implies that the ρ values are strictly upper limits, i.e. another (probably fainter) galaxy may exist at smaller impact parameter; and (ii) there may be little physical connection (e.g. gravitational interaction) between this gas and the galaxy on such large scales. Nevertheless, the results may offer a valuable constraint for galaxy/IGM models and provide further insight into the galaxy/absorber connection.

There are two obvious conclusions to draw from Figure 6: (a) the median equivalent width decreases with increasing ρ ; and (b) the detection rate declines beyond $\rho \approx 300$ kpc but apparently remains above the random rate ($\approx 10\%$; § 3.1) out to at least 1 Mpc. These conclusions are not independent, e.g. the median $W^{\text{Ly}\alpha}$ value is significantly lower at $\rho > 500$ kpc because of the lower detection rate. A Spearman’s correlation test of only

¹³ Our results do not confirm the findings of Wakker & Savage (2009) that galaxies in groups exhibit systematically weaker Ly α absorption, although we have not yet constructed a well-defined group sample from the LCO/WFCCD survey.

TABLE 8
GALAXIES WITHIN 1 MPC OF A QSO SIGHTLINE

Field	RA (J2000)	DEC (J2000)	z_{gal}	ρ (kpc)	L (L_*)	$z_{\text{abs}}^{Ly\alpha}$	$W^{Ly\alpha}$ (mÅ)	N_{HI}	Ref.	$z_{\text{abs}}^{\text{OVI}}$	W^{OVI} (mÅ)	N_{OVI}	Ref
Q0026+1259	00:29:09.32	+13:16:28.5	0.0329	43	0.018	...	...	...	...	0.0329	< 120	< 14.00	9
	00:29:15.36	+13:20:57.0	0.0393	213	1.384	...	...	...	...	0.0393	< 120	< 14.00	9
	00:28:53.55	+13:24:19.9	0.0565	594	0.559	...	...	...	...	...	...	...	...
	00:29:53.44	+13:10:49.5	0.0738	880	0.186	...	...	...	...	...	...	...	...
	00:29:16.48	+13:21:52.9	0.0804	497	0.113	...	...	...	...	...	...	...	...
	00:29:37.55	+13:10:21.0	0.0961	822	1.958	...	...	...	...	0.0961	< 100	< 14.00	9
	00:29:23.49	+13:09:40.8	0.1125	782	1.091	...	...	...	...	0.1125	< 100	< 14.00	9
TONS180	00:29:16.04	+13:10:02.5	0.1314	792	0.179	...	...	...	...	...	...	...	...
	00:57:04.01	-22:26:51.2	0.0234	147	0.019	0.0234	222	13.80	5	0.0234	43	13.48	4
	00:57:08.52	-22:18:29.6	0.0456	265	0.377	0.0456	212	13.80	5	0.0456	62	13.74	4

NOTE. — [The complete version of this table is in the electronic edition of the Journal. The printed edition contains only a sample.]




FIG. 6.— (a) Measurements of associated Ly α absorption for galaxies of all luminosity in our sample within 1 Mpc of a quasar sightline with UV spectral coverage of the Ly α line. We have grouped galaxies together in intervals of 400 km s $^{-1}$ and only plot the system nearest to the sightline. Red diamonds indicate L^* galaxies, blue asterisks are sub- L^* galaxies, and green plus-signs indicate dwarf galaxies. Arrows of the same colors represent non-detections for galaxies of these luminosities. There is a strong anti-correlation between Ly α equivalent width (and column density) and impact parameter. Furthermore, there is a very high incidence of positive detections to 50 mÅ to $\rho \approx 500$ kpc and a marked decline at larger impact parameter. The gray circles and line trace the median equivalent width (including limits) in bins of $\rho = [0 - 100]$ kpc, $[100 - 300]$ kpc, $[300 - 500]$ kpc, $[500 - 1000]$ kpc (Table 6). (b) Detection rate of a Ly α line within 400 km s $^{-1}$ of the galaxies in the LCO/WFCCD survey as a function of impact parameter, for two limiting equivalent widths. Error bars reflect a bootstrap analysis which provide a crude estimate of the uncertainty in the measurements. The dashed red line indicates the prediction for the detection rate for a simplistic filamentary model with filamentary width $w \approx 400$ kpc (see text for details).

the *detections* reports the null hypothesis of no correlation is ruled out at $> 99.99\%$ c.l. This anti-correlation is only strengthened by the non-detections, and it is not a natural consequence of incompleteness in the galaxy survey. We have performed a simple linear regression on the positive detections in log-log space, i.e. by assuming $W^{Ly\alpha}(\rho) = W_0(\rho/1 \text{ kpc})^{-\gamma}$. If we ignore the L^* galaxies (for reasons discussed below) and perform a linear regression on the remaining galaxies with detections, we recover $W_0 = 4.0 \text{ \AA}$ and $\gamma = -0.69$. These values are in good agreement with Tripp et al. (1998), whose analysis was performed on a heterogenous set of galaxy samples. We also note that the $W^{Ly\alpha}$ values have a shallower trend with impact parameter for $\rho > 100$ kpc. Removing those galaxy/absorber pairs from the linear regression, we recover $W_0 = 3.3 \text{ \AA}$ and $\gamma = -0.43$. These values may better describe evolution in the large-scale structures that surround low- z galaxies. We also stress that a rank-correlation test on the detections and upper limits for galaxies at $\rho > 200$ kpc rules out the null hypothesis at $> 99.93\%$ c.l. Both the detection rate and median equivalent width of Ly α absorption are correlated with galaxy impact parameter to at least 1 Mpc, i.e. far beyond the virialized halos and presumed CGM of these galaxies. Before concluding, we note that a power-law description for $W^{Ly\alpha}(\rho)$ was not physically motivated. Furthermore, despite the significant anti-correlation, it is not a very good description of the data; there is tremendous scatter about the regression at all impact parameters.

A closer inspection of Figure 6 reveals that the Ly α detections for L^* galaxies do not follow the trend of decreasing $W^{Ly\alpha}$ with increasing ρ . In fact, the $W^{Ly\alpha}$ values for these galaxies form two sets: the galaxies with positive detections show $W^{Ly\alpha} > 100 \text{ mÅ}$ (generally $W^{Ly\alpha} > 200 \text{ mÅ}$), yet approximately half of the L^* galaxies have no associated absorption to very sensitive limits ($W^{Ly\alpha} < 50 \text{ mÅ}$). This phenomenon is highlighted in Figure 7 where we plot $W^{Ly\alpha}$ vs. ρ for every L^* galaxy within 1 Mpc, showing the closest galaxy to the sightline in cases where two or more L^* galaxies lie within $\pm 400 \text{ km s}^{-1}$ of one another (Table 9). Again, the positive detections primarily result in $W^{Ly\alpha} > 200 \text{ mÅ}$ and approximately half of the systems (without a fainter, neighboring galaxy with smaller ρ) have a non-detection

TABLE 9
 $L > L^*$ GALAXIES WITHIN 1 MPC OF A QSO SIGHTLINE

Field	RA (J2000)	DEC (J2000)	z_{gal}	T ^a	ρ (kpc)	L (L^*)	$\mathcal{N}^b$	$z_{\text{abs}}^{\text{Ly}\alpha}$	$W^{\text{Ly}\alpha}$ (mÅ)	N_{HI}	Rf.	$z_{\text{abs}}^{\text{OVI}}$	W^{OVI} (mÅ)	N_{OVI}	Rf.
Q0026+1259	00:29:15.36	+13:20:57.0	0.0393	E	213	1.384	0	...	...	...		0.0393	< 120	< 14.00	9
	00:29:37.55	+13:10:21.0	0.0961	E	822	1.958	0	...	...	...		0.0961	< 100	< 14.00	9
	00:29:23.49	+13:09:40.8	0.1125	L	782	1.091	0	...	...	...		0.1125	< 100	< 14.00	9
PKS0312-77	03:12:31.11	-76:43:25.0	0.0514	E	692	4.904	3	...	...	...		...	...	...	...
	03:11:02.46	-77:00:05.2	0.0529	E	897	1.477	3	...	...	...		...	...	...	...
	03:11:58.58	-76:48:55.4	0.1192	L	366	1.846	1	0.1183	341	14.17	3	...	...	...	...
PKS0405-123	04:08:06.63	-12:12:50.9	0.0800	E	399	1.404	4	0.0814	254	13.76	3	0.0800	< 40	< 13.50	9
	04:07:54.22	-12:14:50.7	0.0967	E	355	1.357	9	0.0966	484	14.64	3	0.0966	59	13.71	3
	04:07:51.28	-12:11:38.3	0.1670	L	115	2.143	1	0.1671	697	15.47	3	0.1669	412	14.64	1
	04:07:42.79	-12:11:33.1	0.2030	E	262	1.070	9	0.2030	< 50	< 13.00	9	0.2030	< 30	< 13.40	9
	04:08:01.90	-12:11:40.1	0.2484	L	736	1.289	1	0.2484	< 45	< 12.90	9	0.2484	< 30	< 13.40	9
	04:07:34.68	-12:13:22.6	0.2951	E	950	1.198	3	0.2951	< 90	< 13.20	9	0.2951	< 30	< 13.40	9
	04:07:50.69	-12:12:25.2	0.2976	E	245	1.878	7	0.2977	264	13.85	3	0.2976	< 50	< 13.60	9
	04:07:52.33	-12:14:08.0	0.3099	E	689	1.791	3	0.3099	< 40	< 12.90	9	0.3099	< 40	< 13.50	9
	04:07:57.96	-12:09:52.5	0.3203	L	771	1.038	1	0.3203	< 50	< 13.00	9	0.3203	< 30	< 13.40	9
	04:07:50.24	-12:09:52.2	0.3252	E	475	2.317	0	0.3250	188	13.81	3	0.3252	< 50	< 13.60	9
	04:07:45.96	-12:11:09.9	0.3612	E	213	6.604	5	0.3608	554	15.14	3	0.3616	96	14.00	1
	04:07:51.88	-12:13:16.6	0.4253	E	585	6.161	5	...	...	...		...	...	...	...
	04:07:55.50	-12:10:37.0	0.4282	U	638	1.086	2	...	...	...		...	...	...	...
	04:07:41.00	-12:13:15.4	0.5563	U	893	1.854	0	...	...	...		...	...	...	...
PG1004+130	10:07:34.55	+12:52:09.5	0.0704	E	290	2.404	0	...	...	...	...	0.0704	< 100	< 13.90	9
	10:07:25.39	+12:53:04.8	0.1674	E	665	3.577	2	...	...	...		0.1674	< 80	< 13.80	9
	10:07:10.67	+12:50:04.3	0.1931	L	722	1.339	3	...	...	...		0.1931	< 80	< 13.80	9
PG1116+215	11:19:24.31	+21:10:30.6	0.0590	E	614	1.419	3	0.0593	187	13.64	1	0.0593	63	13.77	1
	11:19:06.73	+21:18:29.3	0.1383	E	129	2.039	4	0.1385	479	> 14.35	1	0.1385	83	13.95	1
	11:19:16.88	+21:14:57.1	0.1652	L	761	1.150	14	0.1655	765	> 14.43	1	0.1655	111	14.08	1
PG1211+143	12:14:01.15	+14:11:07.2	0.0522	E	508	1.210	6	0.0510	703	15.67	3	0.0513	...	14.30	4
PG1216+069	12:19:30.89	+06:43:34.6	0.0805	L	470	2.252	1	0.0805	379	13.87	3	0.0815	< 120	< 14.00	9
	12:19:40.12	+06:32:01.5	0.1191	L	983	2.326	3	0.1184	< 30	< 12.70	9	0.1191	< 130	< 14.10	9
	12:19:33.88	+06:42:44.2	0.1250	E	656	1.789	4	0.1236	1433	> 14.78	1	0.1236	378	14.70	1
	12:19:40.27	+06:40:41.2	0.1354	E	706	2.213	4	0.1350	452	15.12	3	0.1354	< 130	< 14.10	9
	12:19:06.47	+06:39:28.8	0.1810	L	632	2.391	3	0.1808	205	14.38	3	0.1810	< 80	< 13.80	9
	12:19:04.05	+06:41:35.6	0.1917	U	919	1.143	3	0.1917	< 45	< 12.90	9	0.1917	< 40	< 13.50	9
	12:19:20.39	+06:36:57.6	0.2464	E	363	1.756	0	0.2464	< 45	< 12.90	9	0.2464	< 40	< 13.50	9
	12:19:34.84	+06:40:55.8	0.2795	L	990	1.832	7	0.2786	49	12.98	6	0.2795	< 60	< 13.60	9
	12:28:51.89	+02:06:03.2	0.0902	E	443	1.105	0	0.0902	161	13.53	1	0.0902	16	13.13	1
	12:29:24.10	+02:08:12.5	0.1464	L	957	2.024	1	0.1466	414	13.94	3	0.1466	< 9	< 12.87	4
PKS1302-102	13:05:20.22	-10:36:30.4	0.0426	L	212	2.577	1	0.0422	410	14.83	3	0.0422	194	14.38	3
	13:05:14.84	-10:40:00.4	0.0569	E	499	1.117	0	0.0569	< 350	< 15.00	9	0.0569	< 40	< 13.50	9
	13:05:41.01	-10:26:42.5	0.0652	E	484	1.946	0	0.0647	256	14.06	3	0.0647	69	13.70	3
	13:05:30.32	-10:26:17.2	0.0718	E	540	2.399	1	0.0712	< 500	< 16.00	9	0.0718	< 40	< 13.50	9
	13:05:20.96	-10:34:51.5	0.0939	E	330	5.381	9	0.0949	681	15.35	3	0.0948	84	13.75	4
	13:05:28.64	-10:38:14.9	0.1384	E	691	1.197	5	0.1393	< 35	< 12.80	9	...	...	...	...
	13:05:40.17	-10:36:54.2	0.1429	E	563	2.051	1	0.1429	< 40	< 12.90	9	0.1429	< 40	< 13.50	9
	13:05:34.00	-10:39:59.6	0.1454	L	952	1.071	3	0.1453	890	15.29	3	0.1453	146	14.16	4
	13:05:31.29	-10:35:42.1	0.1924	E	434	1.533	3	0.1916	401	15.01	3	0.1916	80	13.93	1
	21:54:56.64	-09:18:07.9	0.0517	E	251	1.427	0	0.0515	272	14.08	3	0.0514	< 47	< 13.62	4
FJ2155-0922	21:55:22.71	-09:23:51.8	0.0600	E	357	1.745	6	0.0600	< 30	< 12.70	9	0.0600	< 120	< 14.00	9
	21:55:17.31	-09:17:51.8	0.0734	E	471	3.736	3	0.0736	232	14.64	3	0.0734	< 14	< 13.06	4
	21:54:34.63	-09:16:32.6	0.0785	E	741	1.574	21	0.0777	448	> 14.29	1	0.0777	46	13.61	1
	21:55:00.63	-09:31:28.6	0.0831	E	792	2.979	1	...	...	...	...	0.0831	< 50	< 13.60	9
	21:55:06.53	-09:23:25.2	0.1326	L	212	1.626	2	0.1324	487	> 14.30	1	0.1324	225	14.39	1
	21:54:54.91	-09:23:30.8	0.1764	E	330	2.399	0	0.1765	479	> 14.27	1	0.1765	...	14.44	1
	21:59:08.29	-30:20:54.2	0.0454	U	423	1.142	2	0.0453	87	13.40	3	0.0453	< 7	< 12.76	4
	21:58:23.81	-30:19:31.2	0.0541	E	543	2.053	0	0.0540	315	14.06	3	0.0540	32	13.63	3
	21:58:40.81	-30:19:27.1	0.0570	E	405	1.595	0	0.0566	477	14.48	3	0.0571	44	13.57	3

REFERENCES. — 1: Tripp et al. (2008); 2: Thom & Chen (2008b); 3: Danforth & Shull (2008); 4: Danforth et al. (2006); 5: Penton et al. (2004); 6: Chen & Mulchaey (2009); 9: This paper.

^a Spectral type (E=Early-type, L=Late-type, U=unknown; see text for the quantitative definition).

^b Number of additional galaxies within 3 Mpc of the sightline, 400 km s⁻¹ of this galaxy, and having $L > 0.1L^*$.




FIG. 7.— (a) Measured $\text{Ly}\alpha$ equivalent widths for every L^* galaxy in the LCO/WFCCD survey (with the closest galaxy shown for cases with two or more within 400 km s^{-1}) as a function of impact parameter, extending to 1 Mpc. Galaxies with early-type spectra are marked with open circles and those with late-type spectra are plotted as asterisks. Small/large arrows denote limits for late/early-type spectra. The squares (triangles) indicate cases where an additional sub- L^* (dwarf) galaxy with consistent redshift lies closer to the sightline. The galaxies exhibit two sets of $W^{\text{Ly}\alpha}$ measurements: (i) positive detections with $W^{\text{Ly}\alpha} > 200 \text{ mÅ}$ and (ii) a population of non-detections with $W^{\text{Ly}\alpha} < 100 \text{ mÅ}$ (generally $< 50 \text{ mÅ}$). (b) Equivalent width of O VI 1031 as a function of impact parameter for L^* galaxies from the LCO/WFCCD survey. Symbols have the same meaning as in (a). Only a small fraction of these galaxies exhibit a positive O VI detection for $\rho > 250 \text{ kpc}$, especially if one excludes the systems where a closer, fainter galaxy exists at the same redshift. These results indicate that L^* galaxies are not surrounded by a CGM to many hundred kpc with a high covering fraction to significant O VI absorption.

with $W^{\text{Ly}\alpha} < 100 \text{ mÅ}$. We find no clear differences in the $\text{Ly}\alpha$ detection rate or $W^{\text{Ly}\alpha}$ values for these galaxies as a function of spectral type or redshift. We hypothesize that the subset without a detection could be sightlines penetrating a hot (e.g. virialized) gas which has collisionally ionized hydrogen to non-detectable levels. The positive detections, in contrast, would arise from gas that has not been shock-heated or has since cooled. This simple scenario, however, may not naturally explain the nearly constant $W^{\text{Ly}\alpha}$ values for the positive detections.

To summarize the discussion thus far, our results (and previous work) demonstrate that galaxies of all types are surrounded by a highly ionized medium (which we term as an extended CGM) with near unity covering fraction to $\text{Ly}\alpha$ absorption to $\rho \approx 300 \text{ kpc}$. Furthermore, galaxies of all types exhibit a declining detection rate for H I absorption with $W^{\text{Ly}\alpha} > 50 \text{ mÅ}$ to $\rho \approx 1 \text{ Mpc}$. A key question motivated by these results is what fraction of

the $\text{Ly}\alpha$ forest arises from the CGM surrounding galaxies? More bluntly, can the virialized halos and extended CGM of galaxies account for the majority of lines in the observed $\text{Ly}\alpha$ forest? These questions frame the long-standing debate on whether $\text{Ly}\alpha$ absorbers arise primarily from the gaseous halos of galaxies or from the large-scale, overdense medium (e.g. filaments) that encompass them. To a large degree, the argument hinges on whether the structures giving rise to $\text{Ly}\alpha$ absorption have dimensions of several hundred kpc or are greater than 1 Mpc.

It is evident from the results described above that a fraction of $\text{Ly}\alpha$ absorbers are closely associated with galaxies. Indeed, some fraction of the $\text{Ly}\alpha$ forest must arise from gas bound to individual galaxies. The majority of galaxy/absorber pairs in our sample, however, occur at impact parameters that significantly exceed the presumed virial radius of the galaxies. Indeed, the measured H I column densities are many orders of magnitude smaller than the surface densities characteristic of star-forming galaxies; the gas cannot be related to canonical H I disks nor even the tidal features regularly observed in 21 cm emission observations (e.g. Hibbard et al. 2001). We conclude, therefore, that $\text{Ly}\alpha$ absorption is generally unrelated to the inner galaxy and/or the processes of galaxy formation (e.g. winds, tidal stripping). Instead, the material we have associated with galaxies lies within their extended CGM. In this case the association need not be causal; the gas may not be bound to the galaxy and the two phenomena — galaxies and H I absorbers — may simply trace the same overdensities in the universe.

We draw further insight into the origin of the $\text{Ly}\alpha$ forest from the results presented in Figure 4. As has been emphasized previously (Penton et al. 2002; Chen et al. 2005; Morris & Jannuzi 2006), the origin of $\text{Ly}\alpha$ absorbers appears to be sensitive to the H I column density. At low N_{HI} values, the fraction of ‘random’ absorbers that may be associated to a galaxy with $\rho < 300 \text{ kpc}$ is small ($< 20\%$ [15/82] for $N_{\text{HI}} = 10^{13-14} \text{ cm}^{-2}$). This suggests that the majority of weak $\text{Ly}\alpha$ absorbers, which dominate the $\text{Ly}\alpha$ forest by number, arise beyond the virialized halos and the CGM of individual galaxies. Presumably, these absorbers arise in overdense regions of the low- z universe which have not supported the formation of a luminous galaxy. In contrast, the percentage of strong absorbers ($N_{\text{HI}} > 10^{15} \text{ cm}^{-2}$) with an associated galaxy at $\rho < 300 \text{ kpc}$ is 80% (8/10). One is compelled, for this subset, to physically associate the absorbers with the galaxies. These inferences are supported by the N_{HI} dependence measured for the clustering of $\text{Ly}\alpha$ absorbers to low- z galaxies (Penton et al. 2002; Chen et al. 2005; Chen & Mulchaey 2009; Shone et al. 2010). The much lower cross-correlation amplitude for systems with $N_{\text{HI}} < 10^{14} \text{ cm}^{-2}$ imply this gas is not physically associated to galaxies. The high amplitude for $N_{\text{HI}} > 10^{14.5} \text{ cm}^{-2}$ absorbers, meanwhile, implies a physical association with galaxies. These strong absorbers are sufficiently rare that one may associate each of them with a galaxy. Indeed, the results in Figure 4 suggest that many of the strong absorbers arise within the virialized halos of individual galaxies. Such environments may be the only regions of the universe with sufficient overdensity to give rise to H I column densities exceeding 10^{15} cm^{-2} .

Another statistical approach to addressing these issues is to compare the observed incidence of $\text{Ly}\alpha$ absorp-




FIG. 8.— The horizontal, hatched regions show measurements for the incidence of Ly α at $z = 0$ as a function of limiting equivalent width (Penton et al. 2004). For strong absorbers ($W^{\text{Ly}\alpha} > 300 \text{ m}\text{\AA}$), we observe ≈ 10 lines per unit pathlength (and redshift). The curves represent estimates for $\ell_{\text{Ly}\alpha}(X)$ derived from the virialized halos of galaxies (black, solid) and the extended CGM surrounding galaxies (blue, dotted) integrated from a limiting luminosity. The results indicate that the virialized halos of galaxies account only for the incidence of strong Ly α lines observed in the low- z IGM. Even allowing for the extended CGM around each galaxy (with $r_{\text{CGM}} = 300 \text{ kpc}$), the majority Ly α absorption in the low- z IGM cannot arise from gas surrounding galaxies.

tion against that predicted for gas surrounding galaxies based on our results and simplified assumptions about the virial radii of galaxies (Equation 1). Specifically, we ask whether galaxy halos occupy enough area to account for most Ly α absorbers to a given equivalent width limit. To this end, consider the incidence per unit pathlength dX of Ly α absorption $\ell_{\text{Ly}\alpha}(X)$, where

$$dX = \frac{H_0}{H(z)} (1+z)^2 dz \quad . \quad (4)$$

At $z = 0$, $dX = dz$ and $\ell(X)$ may be evaluated directly from the observed incidence of absorbers per unit redshift¹⁴, $\ell(z)$. The quantity $\ell(X)$ is defined to remain constant in time if the product of the comoving number density n_c of objects giving rise to the absorption and the physical size A_p of the sources remains constant, i.e.

$$\ell(X) = \frac{c}{H_0} n_c A_p \quad . \quad (5)$$

This evaluation assumes that 100% of the area A_p produces detectable absorption. One could include a fac-

tor in the formalism that allows for a non-unity covering fraction. Alternatively, one may consider A_p to be the effective area of the object to absorption.

In Figure 8, hatched, colored bands show current estimates for $\ell_{\text{Ly}\alpha}(X)$ using results from $z = 0$ surveys of Ly α absorbers (Penton et al. 2004; Wakker & Savage 2009). Each band corresponds to a limiting equivalent width, which for low $W^{\text{Ly}\alpha}$ values translates directly to a limiting column density.¹⁵ Overplotted on Figure 8 is a black, solid curve that shows the cumulative inferred contribution to $\ell_{\text{Ly}\alpha}(X)$ of virialized galactic halos as a function of galaxy luminosity. For this calculation, we have estimated n_c from the galaxy luminosity function of Blanton et al. (2003) taking $h = 0.72$. We have also assumed that $A_p^{\text{Ly}\alpha} = \pi r_{\text{vir}}^2$ with r_{vir} given by equation 1. It is obvious that bright galaxies ($L > L^*$) can only contribute a small fraction of the total incidence of Ly α absorbers. Integrating down to $L = 0.01 L^*$, the approximate limit of the LCO/WFCCD survey, we infer $\ell_{\text{Ly}\alpha}^{\text{vir}}(X; L > 0.01 L^*) \approx 15$. Tumlinson & Fang (2005) recovered a similar estimate for galactic halos (in the

¹⁴ Commonly referred to as $n(z)$ or dN/dz . Similarly $\ell(X)$ is often denoted dN/dX .

¹⁵ $W^{\text{Ly}\alpha} = 50 \text{ m}\text{\AA}$ roughly corresponds to $N_{\text{HI}} = 10^{13} \text{ cm}^{-2}$.

context of O VI absorption) in an analysis that also accounted for the clustering of galaxies. The estimate lies below all of the $\ell_{\text{Ly}\alpha}(X)$ estimates shown in Figure 8, except the strongest Ly α lines ($W^{\text{Ly}\alpha} > 300 \text{ m}\text{\AA}$). We conclude, therefore, that *the virialized halos of individual galaxies can only account for the strongest Ly α absorbers ($W^{\text{Ly}\alpha} > 300 \text{ m}\text{\AA}$) observed in the low- z IGM.* On the other hand, the results presented in Figures 1–3 indicate systems with $\rho < r_{\text{vir}}$ to $L > 0.1L^*$ galaxies do exhibit a high covering fraction to $W^{\text{Ly}\alpha} > 300 \text{ m}\text{\AA}$ absorption. One further concludes that *the majority of strong Ly α absorbers must arise from gas in galactic halos.*

As emphasized throughout this section, galaxies are also surrounded by a CGM with nearly unit covering fraction for $W^{\text{Ly}\alpha} > 50 \text{ m}\text{\AA}$ to $\rho = 300 \text{ kpc}$ (Figure 6b). The dashed blue curve in Figure 8 shows an estimate of the cumulative incidence of Ly α absorbers derived under the assumption that a CGM with $r_{\text{CGM}} = 300 \text{ kpc}$ surrounds every galaxy with unit covering fraction, i.e. $A_p^{\text{Ly}\alpha} = \pi r_{\text{CGM}}^2$, independent of galaxy luminosity. This increases the estimated $\ell_{\text{Ly}\alpha}(X)$ value relative to the estimate for virialized halos, especially for galaxies with $r_{\text{vir}} \ll 300 \text{ kpc}$. Nevertheless, even the gas from the extended CGM of galaxies cannot match the observed incidence of weak Ly α absorbers.¹⁶ We conclude that *the overwhelming majority of Ly α lines detected from the IGM arise in structures located at distances beyond several hundred kpc from $L > 0.01L^*$ galaxies.* One must seek an additional origin for the majority of IGM absorption then gas surrounding galaxies.

For nearly two decades, researchers have performed hydrodynamic cosmological simulations of the IGM to simulate and study the Ly α forest (e.g. Miralda-Escudé et al. 1996; Gnedin & Hui 1998). This includes several analyses of the low- z IGM (Davé et al. 1999; Cen & Ostriker 1999; Richter et al. 2006; Paschos et al. 2009). Regarding Ly α absorption, the low- z studies have focused primarily on the distributions of H I column density and Doppler parameter and on associating the gas to various phases of the universe. In their recent publication, Davé et al. (2010) have reported that the majority of observable Ly α absorption (systems with $N_{\text{HI}} \approx 10^{13}\text{--}10^{15} \text{ cm}^{-2}$) is associated with diffuse, photoionized gas at temperatures of $\sim 10^4 \text{ K}$, with density $n_{\text{H}} \approx 10^{-4}\text{--}10^{-6} \text{ cm}^{-3}$, and arising in structures with characteristic sizes of several hundred kpc to 1 Mpc. These structures, which trace large-scale overdensities in the low- z universe, presumably also contain galaxies. In their first paper, Davé et al. (1999) examined the association of galaxies to the IGM and found that the cosmological simulations reasonably reproduced the observed high incidence of Ly α absorption at small impact parameters to galaxies. They also roughly reproduced the observed relation between equivalent width and impact parameter to $\rho \approx 1 \text{ Mpc}$. The majority of these Ly α absorbers arise in shocked or diffuse gas surrounding the galaxies and the gas is not bound to them. The overall incidence of Ly α , meanwhile, is dominated by gas at $\rho \gg 100 \text{ kpc}$ from galaxies. Their results

appear, at least qualitatively, to provide a reasonable match to our new observations although we encourage future, quantitative comparison. We are further compelled, therefore, to support this model over ones where virialized halos or the extended CGM associated with discrete galaxies dominate the observed IGM.

The cosmological simulations described above predict that the gas arises in a filamentary network of overdensities, known as the “cosmic web”, that permeates the universe. In this same framework, galaxies form within the filaments and galaxy clusters are produced at the nodes where filaments intersect. The observed distribution of galaxies in wide-field surveys generally support this picture (Gott et al. 2009; Bond et al. 2010). Inspired by this paradigm, we introduce a simple filament model to interpret the trends observed in Figure 6. The model assumes that each galaxy in the LCO/WFCCD survey resides at the center of a filamentary structure, generally not at a node. The filaments have a finite width w but are presumed to be infinitely long.¹⁷ In this simple scenario, the covering fraction to Ly α absorption associated with a galaxy is purely geometrical. It is unity for $\rho < w/2$ and decreases at larger values. Overplotted in Figure 6b is the predicted covering fraction for this simple model with $w = 400 \text{ kpc}$. This model shows good agreement with the observations despite its simplicity. It suggests that the overdense structures hosting $z \sim 0$ galaxies have characteristic dimension of several hundred kpc. Indeed, Davé et al. (2010) estimate that $w \sim 400 \text{ kpc}$ corresponds to absorbers with $N_{\text{HI}} \approx 10^{13.3} \text{ cm}^{-2}$, a characteristic value for the gas analyzed here. Lastly, if such filaments dominate the incidence of Ly α in the observed IGM then their filling factor of galaxies must be low so that galaxies are only rarely associated with Ly α absorption (Figures 4 and 8). We infer that only rare ‘patches’ of filaments contain $L > 0.01L^*$ galaxies.

4.2. O VI

Let us now consider O VI gas and its relation to galaxies in the $z \sim 0$ universe. Aside from the H I Lyman series, the most commonly detected absorption lines of the low- z IGM are the O VI doublet. This reflects the high number abundance of oxygen, the ease of identifying a line doublet, and the predominance of regions within the IGM that can produce the O^{+5} ion. With the launch of *HST* and its suite of UV spectrometers, O VI has been surveyed from $z \approx 0.2\text{--}1$ (Burles & Tytler 1996; Tripp et al. 2000, 2008; Thom & Chen 2008b; Danforth & Shull 2008). Secondly, the *FUSE* spectrometer, which operated at $\lambda \approx 910\text{--}1150 \text{\AA}$ for nearly a decade, enabled the search for intergalactic O VI down to $z = 0$ (Prochaska et al. 2004; Danforth et al. 2006; Wakker & Savage 2009). Together, these surveys have discovered over 100 O VI doublets to establish their incidence as a function of equivalent width (and column density), to characterize their relation to H I gas, and to measure their distribution of line widths.

Previous work has proposed a variety of origins for this O VI absorption. On the smallest scales, the high incidence of O VI detections in the halo of our Galaxy (Sembach et al. 2003) implies a high filling factor of such

¹⁶ Our estimate could be increased by a modest factor if we allowed that one galaxy may give rise to multiple Ly α absorbers. On the other hand, Bowen et al. (2002) have noted that Ly α ‘clusters’ are likely associated with galaxy groups (i.e. multiple galaxies).

¹⁷ These filaments also have a finite depth whose dimension is irrelevant to the analysis.

gas in the dark matter halos of $L \approx L^*$ galaxies. This assertion has now been confirmed, at least for blue and star-forming L^* galaxies at $z \sim 0.2$ (Tumlinson et al., in prep.). At the largest scales (i.e. intergalactic), cosmological simulations predict that the WHIM comprises $\gtrsim 25\%$ of the baryons in the modern universe, and some models also predict that this gas gives rise to a significant fraction of the observed O VI absorption (e.g. Cen & Fang 2006). In between these extremes, the intragroup medium (revealed in some cases by X-ray emission) may have the density and temperature required to yield significant O VI absorption (Mulchaey et al. 1996). Of course, each of these environments is likely to contribute, and each could even dominate at different O VI equivalent width.

A key approach to discriminating among these scenarios is to explore the association between O VI gas and galaxies. Tumlinson & Fang (2005) compared the incidence of O VI absorption against predictions for low- z galaxies as characterized by SDSS and concluded that bright galaxies ($L > L^*$) could not reproduce the observed rate. They suggested that fainter galaxies may explain the observed incidence if dwarf galaxies were surrounded by an enriched medium to ≈ 200 kpc. Stocke et al. (2006) have correlated $L > 0.1L^*$ galaxies with 9 O VI absorbers in several fields and reported a median offset of $\approx 180 h_{70}^{-1}$ kpc. They argued that O VI gas is predominantly associated with individual galaxies and speculated that galaxies with $L < 0.1L^*$ dominate. Wakker & Savage (2009) surveyed 6 field galaxies at $z \approx 0$ with $L \geq 0.1L^*$ and $\rho < 350$ kpc and reported O VI detections for 2/3. They also argued that gas in the extended surroundings of galaxies contributes significantly to O VI (and Ly α) absorption.

The results presented in this paper offer new insight into the origin of O VI at modest to large equivalent widths ($W^{1031} \gtrsim 30$ mÅ). Our survey and analysis has focused on the properties of galaxy halos and the extended CGM as related to Ly α and O VI absorption. In contrast to Ly α (as discussed in the previous subsection), the association between galaxies and O VI shows a strong dependence on galaxy luminosity. In § 3.1 (Figure 1), we showed that sightlines at impact parameters $\rho < 300$ kpc to dwarf galaxies ($L < 0.1L^*$) rarely exhibit O VI absorption. Furthermore, the few dwarfs with associated O VI gas also had a neighboring $L > 0.1L^*$ galaxy within $\rho = 200$ kpc of the sightline. The implication is that the extended CGM of dwarf galaxies¹⁸ does not contribute significantly to O VI absorption with $W^{1031} \gtrsim 50$ mÅ. For the brightest galaxies ($L > L^*$), we found a high incidence of O VI detections for $\rho < 225$ kpc but not a single detection beyond (to sensitive limits; Figure 2, Table 4). In Figure 7b, we have extended the search for O VI associations with L^* galaxies to 1 Mpc. Although there are a few detections for $\rho > 300$ kpc, the majority of these cases also show an additional, fainter ($L < L^*$) galaxy located within $\rho = 300$ kpc of the sightline. We conclude that the covering fraction of O VI gas around L^* galaxies is small for $\rho \gtrsim 250$ kpc (see also Wakker & Savage 2009).

In contrast to the faint and bright galaxies, the inter-

mediate sub- L^* population exhibits a high incidence of O VI at all impact parameters $\rho < 300$ kpc (Figure 3). This conclusion is independent of spectral type; we find associated O VI for sub- L^* galaxies with early-type (presumed ‘red and dead’) and late-type (star-forming) spectra. Combining these results, we associate O VI gas preferentially with the halos and CGM of sub- L^* galaxies. But this begs the obvious question, are these galaxies sufficiently common that they trace the majority of O VI in the universe? Or does O VI also arise from an additional reservoir, as we found for weak Ly α absorbers?

In Figure 9, we plot the incidence per unit path-length dX of O VI absorption, $\ell_{\text{OVI}}(X)$, estimated from the incidence per unit redshift $\ell_{\text{OVI}}(z)$ measurements of Tripp et al. (2008) for several equivalent width limits. Formally, we have converted their estimates of the incidence per unit redshift $\ell(z)$ measured at $z \approx 0.2$ to $\ell(X)$ (Equation 4), which implies a downward correction of $\approx 30\%$. The width of the hatched regions indicate the 1σ uncertainties reported by Tripp et al. (2008) and we caution that systematic bias may have led to a modest overestimate in these $\ell_{\text{OVI}}(X)$ evaluations (Thom & Chen 2008a). Overlaid on the observational estimates of $\ell_{\text{OVI}}(X)$, we plot the predicted incidences of O VI from each of our galaxy subsets. The solid bars show the estimated contribution from gas within the virialized halos of each galaxy sample. In each case, we have estimated n_c from the galaxy luminosity function¹⁹ of Blanton et al. (2003) taking $h = 0.72$. For the effective cross-section of halos, we have assumed $A_p^{\text{OVI}} = \pi r_{\text{vir}}^2$ with $r_{\text{vir}} = 100, 160, 250$ kpc for the dwarf, sub- L^* , and L^* populations, respectively. These values are somewhat smaller than the value one would derive by averaging Equation 1 over the luminosity function as our estimates for A_p^{OVI} include a modest correction for the covering fractions to O VI absorption.

None of the galaxy populations (on their own) can account for the incidence of the commonly observed O VI absorbers if one restricts to virialized halos. For the L^* galaxies, the estimated incidence is $\approx 10\%$ of the total, in line with the model predictions by Tumlinson & Fang (2005). Furthermore, dwarf galaxies may give rise to as much as 20–30% of the total incidence but we caution that the estimate shown in Figure 9 assumes a higher covering fraction to O VI for dwarfs than supported by our observations (Figure 1). Together, galactic halos can account for the incidence of very strong O VI absorption ($W^{1031} > 100$ mÅ) and possibly moderate O VI absorbers ($W^{1031} \approx 70$ mÅ). We conclude, however, that *gas in the virialized halos of $z \sim 0$ galaxies with $L > 0.01L^*$ cannot account for the majority of observed O VI systems* ($W^{1031} > 30$ mÅ). A similar inference was drawn by Tumlinson & Fang (2005).

The above conclusion is also supported by the results presented in Figure 5, where we found that very few O VI absorbers are coincident with the virialized halo of a galaxy. We further emphasize that the $z \sim 0$ luminosity function is too shallow ($\Phi \propto L^{-1}$) for fainter galaxies to qualitatively modify this conclusion.²⁰ On

¹⁸ Gas within the virialized halos of dwarf galaxies (i.e., at $\rho \lesssim 50$ kpc), however, was not well probed by our survey.

¹⁹ For the dwarf galaxy calculation, we limit the integration to $0.01L^* < L < 0.1L^*$.

²⁰ Although there remains debate on the precise slope of the




FIG. 9.— Horizontal bands show the observed incidence of O VI absorption per absorption length $\ell_{\text{OVI}}(X)$, as a function of limiting equivalent width (Tripp et al. 2008). The solid vertical bars indicate estimates of the incidence of O VI absorption for gas in the virialized halos of dwarf (black; $0.01L^* < L < 0.1L^*$), sub- L^* (blue; $0.1L^* < L < L^*$), and $L > L^*$ (red) galaxies. For these estimates we have assumed virial radii of $r_{\text{vir}} = 100, 160$ and 250 kpc respectively, and that the halos have a covering fraction of 100% to O VI. Under this parameterization, virialized halos of galaxies only account for stronger O VI systems. The hatched, blue bar gives the estimate of $\ell_{\text{OVI}}(X)$ for gas in the extended CGM of sub- L^* galaxies assuming a 100% covering fraction to $\rho = r_{\text{CGM}} = 300$ kpc, as suggested by our LCO/WFCCD survey (Figure 3). The excellent correspondence with the observed $\ell_{\text{OVI}}(X)$ value for $W^{1031} > 30$ mÅ indicates that the remainder of O VI systems arise in this extended CGM. We conclude that the majority of O VI gas observed in the low- z IGM arises in the diffuse medium surrounding individual galaxies (predominantly $L \approx 0.3L^*$), and that it rarely originates in the WHIM predicted by cosmological simulations.

the other hand, one may attribute the strongest O VI absorbers ($W^{1031} > 100$ mÅ) to virialized halos and, by inference, galactic-scale processes. In fact, one must conclude from the results presented in Figures 1-3, 5, and 9 that *the majority of strong O VI systems arise from within the halos of individual galaxies.*

With virialized halos ruled out as the dominant origin of O VI gas (by frequency of absorption not necessarily by mass), we now consider the extended CGM surrounding $z \sim 0$ galaxies. As emphasized above, we find a significant covering fraction to $\rho = 300$ kpc for the sub- L^* galaxies (Figure 3). Overlaid on the $\ell_{\text{OVI}}(X)$ estimate for virialized halos of the sub- L^* galaxies is a hatched bar that represents the contribution to $\ell_{\text{OVI}}(X)$ from an extended CGM surrounding each sub- L^* galaxy. This hatched bar assumes $A_p^{\text{OVI}} = \pi r_{\text{CGM}}^2$ with $r_{\text{CGM}} = 300$ kpc and a 100% covering fraction for $W^{1031} \geq 30$ mÅ. This amounts to a several times higher

faint-end, one would require a much steeper slope than currently estimated for faint galaxies to significantly contribute.

incidence of O VI absorption. Remarkably, we find that the extended CGM of sub- L^* galaxies reproduces the incidence of all O VI absorbers with $W^{1031} \gtrsim 30$ mÅ! Of course, these results follow from the close association of galaxies to O VI absorbers (Figure 5) and the high covering fraction of O VI gas surrounding sub- L^* galaxies (Figure 3). While one could reduce this estimate by a factor of order unity (e.g. by assuming a smaller covering fraction), *we conclude that the extended CGM of sub- L^* galaxies is the primary reservoir of O VI gas observed in the $z \sim 0$ IGM.* This final conclusion, which draws a distinction between virialized halos of sub- L^* galaxies and their extended CGM, is sensitive to the precise definition of a galaxy halo and its radius. We encourage additional theoretical exploration into the nature and extent of virialized gas in the halos of subluminal, low- z galaxies.

Let us now consider the implications of an extended CGM that gives rise to significant O VI absorption surrounding sub- L^* galaxies. First, we note that this CGM may include neighboring galaxies, the intragroup medium, and the diffuse medium that envelops the galax-

ies (e.g. filaments). Regarding neighboring galaxies,²¹ we have already demonstrated that virialized halos contribute only modestly to the incidence of O VI. By the same argument, they cannot dominate the detections associated with the extended CGM of sub- L^* galaxies. Regarding the intragroup medium, only a subset of sub- L^* galaxies exhibit galaxy overdensities suggestive of a group (Table 5). We infer, therefore, that this extended CGM is best described by a diffuse, modestly overdense ($\delta \gtrsim 10$) medium. In the IGM paradigm of undulating Gunn-Peterson absorption, such overdensities naturally give rise to significant H I absorption. Indeed, as discussed above, Ly α absorption is a ubiquitous phenomenon for galaxies of all luminosity, and every O VI absorber detected to date shows corresponding Ly α absorption (Thom & Chen 2008b).

Although one may naturally expect H I absorption from an overdense region of the universe, the detection of O VI further requires a chemically enriched medium and the physical conditions that produce the O⁺ ion. Regarding the former, our results require that the CGM of sub- L^* has been polluted by metals on scales of several hundred kpc. Although many of the sub- L^* galaxies in our sample show the spectral signatures of ongoing star-formation, these are not star-bursting systems, and they are unlikely to currently be driving galactic-scale winds to such large distances. We infer, therefore, that this CGM was previously enriched. Current estimates for the metallicities of the O VI absorbers suggest values ranging from 0.01 to nearly solar abundance with a median of ≈ 0.1 solar (e.g. Prochaska et al. 2004; Cooksey et al. 2008; Thom & Chen 2008a; Danforth & Shull 2008). This enrichment level exceeds predictions from the first stars (e.g. Wise et al. 2010) indicating a subsequent phase of star-formation is required. The medium is too diffuse to form metals *in situ*, and therefore, these must have been transported by distances of 100 kpc or more. The proximity of the sub- L^* galaxy suggests that the oxygen was produced in that system, yet this is purely speculative. We also note, following the mass estimate from § 4.1 (Equation 3), that a 0.1 solar metallicity implies a metal mass equivalent to $\approx 3 \times 10^9 M_\odot$ of solar metallicity stars. This could exceed the total mass in metals within a typical sub- L^* galaxy and its satellites, which would require mass outflow rates from galactic winds that match or exceed the star-formation rate (e.g. Weiner et al. 2009). Comprehensive modeling of the chemical enrichment of the IGM that includes galaxy formation and metal transport are required to further explore this topic (e.g. Cen & Ostriker 2006; Ganguly et al. 2008; Oppenheimer & Davé 2008, 2009).

In addition to containing oxygen, this CGM must have the appropriate density and/or temperature for a significant fraction of oxygen to exist as O⁺. One process to produce O⁺ is via collisional ionization (CI) which requires temperatures on the order of $T \sim 10^5$ – 10^6 K. The other obvious mechanism to produce O⁺ is with photoionization, which requires photons with energies $h\nu > 8$ Ryd. Such hard photons are only produced by AGN, and the extragalactic UV background (EUVB) is the obvious source of such radiation when one is far from

active galaxies.

Consider first several inferences one may draw from our association of O VI to the extended CGM of sub- L^* galaxies. If the extended CGM (almost by definition) lies beyond the virialized radius of these galaxies, one may not expect a collisionally ionized medium. Furthermore, the majority of these sub- L^* galaxies are not obviously located within a larger dark matter halo (e.g. a group or cluster). Therefore, this gas may not have been virialized to $T \gg 10^4$ K. One would require instead gravitationally-induced shock-heating from larger scales. Indeed, the collapse of large-scale density ‘waves’ is predicted to drive the production of the WHIM in cosmological simulations (e.g. Cen & Ostriker 1999; Davé et al. 2001) and may, in principle, heat the CGM of sub- L^* galaxies. Although this is plausible, these waves are generally predicted to ‘crash’ on larger scales (many Mpc) than the several hundred kpc scales of the CGM. Indeed, Ganguly et al. (2008) analyzed the average impact parameter to galaxies of O VI gas arising in the WHIM. They found in their simulations that only $\approx 1\%$ of O VI detections with $W^{1031} \geq 50$ mÅ would have a $L > 0.01L^*$ galaxy within 300 kpc. This prediction is strongly ruled out by our observations (see also Wakker & Savage 2009). By inference, we conclude that *O VI gas observed in the low- z IGM does not typically trace a collisionally ionized WHIM.*

Empirically, the ionization mechanism for observed O VI absorbers remains a matter of great debate. We support (and have contributed to) arguments that conclude most O VI systems are not collisionally ionized (Prochaska et al. 2004; Tripp et al. 2008; Thom & Chen 2008a). The evidence includes the close alignment of H I gas with O VI, the narrowness of the coincident Ly α absorption (and of O VI itself), and the very frequent detection of coincident C III absorption. On the other hand, see Fox (2011) for a set of arguments against O VI arising in predominantly photoionized gas. Given the results presented above, we may at least consider whether a photoionized, extended CGM of sub- L^* galaxies may naturally support O VI absorption.

Under the assumptions that the gas is optically thin to ionizing radiation and that the UV radiation field is dominated by a typical AGN spectrum ($f_\nu \propto \nu^{-1.6}$), standard photoionization calculations indicate that an ionization parameter²² $U \gtrsim 10^{-1}$ is required to produce a significant fraction of O⁺ ions. To achieve $U \gtrsim 10^{-1}$ with current estimates for the EUVB (e.g. CUBA, which predicts $J_\nu \approx 10^{-25}$ erg s⁻¹ Hz⁻¹ cm⁻² at 8 Ryd; Haardt & Madau 1996), this demands a volume density $n_H \lesssim 10^{-5}$ cm⁻³. This value is roughly 50 $\times$ the mean baryonic density at $z \sim 0$ suggesting an overdensity $\delta \lesssim 50$. These same photoionization calculations imply a size for the structure of $d \sim 300$ kpc ($N_{HI}/10^{14}$ cm⁻²). The constraints on n_H and d give plausible values for the extended CGM of sub- L^* galaxies. They imply a gas that is overdense yet not virialized ($\delta \ll 200$). Furthermore, the inferred size of the structures coincides with the dimensions inferred from our analysis. We conclude that the gas in the extended CGM could be predominantly photoionized material giving rise to O VI absorption.

²¹ Also, by definition the volume beyond r_{vir} must have a lower density than the medium within it.

²² $U \equiv \Phi/cn_H$, where Φ is the flux of ionizing photons.

We have developed the following picture for O VI absorption in the low- z IGM: the gas has density $n_H \approx 10^{-5} \text{ cm}^{-3}$, is predominantly photoionized, and is primarily located in the extended CGM of sub- L^* galaxies. The oxygen was produced in a previous episode of star-formation (perhaps by the observed sub- L^* galaxy) and has been transported to the CGM by one or more processes. Remarkably, this scenario is qualitatively consistent with the model for O VI described by Oppenheimer & Davé (2009). In their cosmological simulations, oxygen is produced at earlier times (often $z > 1$) in a star-bursting galaxy whose galactic-scale wind transports the metals to the surrounding, overdense IGM. This gas is observed today primarily as photoionized O VI and is coupled to significant H I absorption mainly because both exist in overdense regions.

Oppenheimer & Davé (2009) studied the equivalent width and Doppler parameter distribution of O VI gas and found their model gives reasonably good agreement to the observed distribution²³. These authors also performed a qualitative analysis on the connection between O VI absorption and galaxies (their Figures 14 and 15). They found O VI gas is generally located at several virial radii ($\sim 100\text{--}300 \text{ kpc}$) from galaxies with masses $M_{\text{gal}} \sim 10^{9.5\text{--}10} M_{\odot}$ ($\sim 0.03\text{--}0.1 M^*$). Although these separations are consistent with our results, the typical galaxy mass may be lower than that of the average sub- L^* galaxy. We encourage further analysis to consider the covering fraction to O VI absorption as a function of galaxy luminosity (e.g. Ganguly et al. 2008). The data also permits one to examine trends between the galactic mass and environment with the observed equivalent widths of absorption.

Before concluding, it is worth speculating on why the extended CGM for L^* and dwarf galaxies would have a lower covering fraction than that observed for sub- L^* galaxies. Regarding dwarf galaxies, there are several plausible explanations. First, these galaxies generally show lower N_{HI} values on average suggesting lower total gas columns. Second, it is possible that the CGM of dwarf galaxies is chemically poor in comparison to brighter galaxies.²⁴ Lastly, the physical conditions of the gas (temperature, density) in the CGM surrounding dwarf galaxies may not support the O^{+5} ion, e.g. the gas may have too low volume density. Together, these effects could reduce the associated O VI absorption to equivalent widths below typical detection limits.

For the L^* galaxies, however, we observe H I column densities at least as large as those observed for the sub- L^* galaxies (Table 6). One may also expect the gas to have higher metallicity.²⁵ If the lower incidence of O VI absorption is unrelated to differences in chemical enrichment, then the result must relate to the physical conditions of the gas, i.e., the extended CGM of L^* galaxies

has a density and/or temperature that is not favorable for O^{+5} ions. Under the assumption of photoionization by the EUVB, the gas must have $n_H \approx 10^{-5} \text{ cm}^{-3}$ for its ionization potential to favor O^{+5} ions. If the CGM of L^* galaxies is $\approx 10\times$ higher than that of sub- L^* galaxies at $\rho \sim 200 \text{ kpc}$, then this could explain the preponderance of non-detections. In this scenario, one would predict the detection of metals at $\rho \sim 200 \text{ kpc}$ from L^* galaxies in lower ionization states (e.g. C IV, C III). One might then expect a higher incidence of detections at somewhat larger impact parameters,²⁶ but this is not observed (Figure 7b). As a final alternative, it is possible that gas at $\rho \sim 250 \text{ kpc}$ from an L^* galaxy has been shock-heated to $T \gg 10^5 \text{ K}$. Indeed, these galaxies are more frequently members of a group, i.e. embedded within a larger, virialized halo.

5. SUMMARY

With this paper we have examined the associations of galaxies to $\text{Ly}\alpha$ and O VI absorption in the low- z IGM to explore the origin of this gas. Specifically, we have correlated galaxies from our LCO/WFCCD galaxy survey (Prochaska et al. 2011) and IGM absorption from published line-lists of *HST* and *FUSE* UV quasar spectra. These two datasets are essentially independent of one another, although we have supplemented the IGM line-lists with our own analysis of the quasar sightlines. The following bullets summarize our main findings:

- Galaxies of all luminosity ($L > 0.01 L^*$) and spectral-type show strong, associated $\text{Ly}\alpha$ absorption to impact parameter $\rho \approx 300 \text{ kpc}$ with a very high covering fraction ($\approx 90\%$). The strongest H I absorbers ($W^{\text{Ly}\alpha} \gtrsim 1 \text{ \AA}$) are preferentially associated with brighter galaxies ($L > 0.1 L^*$).
- We estimate a baryonic mass for this extended circumgalactic medium (CGM) of $M_{\text{CGM}} \approx 3 \times 10^{10} M_{\odot}$ ($(N_{\text{H,CGM}}/10^{19} \text{ cm}^{-2})(r_{\text{CGM}}/300 \text{ kpc})^2$), having adopted a constant H I column density $N_{\text{H,CGM}}$.
- Galaxies with luminosities $L > 0.1 L^*$ exhibit a high covering fraction ($> 80\%$) for significant O VI absorption to $\rho = 200 \text{ kpc}$ and to 300 kpc for sub- L^* galaxies. Dwarf galaxies ($L < 0.1 L^*$) exhibit a low covering fraction for $\rho > 50 \text{ kpc}$.
- Despite these high covering fractions of $\text{Ly}\alpha$ and O VI absorption, there are examples of non-detections to very sensitive limits, even at very low impact parameters ($\rho \lesssim 50 \text{ kpc}$).
- We observe a declining covering fraction and median equivalent width for $\text{Ly}\alpha$ to $\rho = 1 \text{ Mpc}$. The equivalent widths of the positive detections at $\rho > 100 \text{ kpc}$ may be described by a power-law, $W^{\text{Ly}\alpha}(\rho) = 3.3 \text{ \AA} (\rho/1 \text{ kpc})^{-0.43}$, but with great scatter.
- To $\rho = 1 \text{ Mpc}$, L^* galaxies show two distinct distributions of $\text{Ly}\alpha$ absorption: (i) a set of positive

²³ These authors had to include turbulent broadening, added in post-processing, to explain the observed linewidths of O VI.

²⁴ An ongoing Cycle 18 *HST*/COS program will assess the chemical enrichment of gas associated with dwarf galaxies (PI: Tumlinson; 1224).

²⁵ Although, one could also speculate that the larger potential well of L^* galaxies prevents the transport of metals beyond their virialized halos (e.g. Oppenheimer & Davé 2008). This hypothesis, however, is apparently not supported by current research on galactic-scale winds (e.g. Weiner et al. 2009; Rubin et al. 2010).

²⁶ Also, low column densities at all impact parameters if one assumes spherical symmetry.

detections with $W^{\text{Ly}\alpha} > 200 \text{ m}\text{\AA}$; and (ii) a set of non-detections with $W^{\text{Ly}\alpha} < 50 \text{ m}\text{\AA}$.

- The detection rate of Ly α absorption with impact parameter may be described by galaxies within filaments having characteristic widths $w \sim 400 \text{ kpc}$.
- Few, if any, of the weak Ly α absorbers ($W^{\text{Ly}\alpha} < 100 \text{ m}\text{\AA}$) from the low- z IGM arise in the virialized halos of $z \sim 0$ galaxies, or their surrounding CGM. The majority of strong Ly α absorbers ($W^{\text{Ly}\alpha} > 300 \text{ m}\text{\AA}$), however, does arise in these environments.
- The strongest O VI absorbers ($W^{1031} > 100 \text{ m}\text{\AA}$; $N_{\text{OVI}} > 10^{14} \text{ cm}^{-2}$) arise preferentially in the galactic halos of $L > 0.01 L^*$ galaxies. Weaker O VI

absorbers are associated with the extended CGM of sub- L^* galaxies.

- Current predictions for models where O VI gas arises in a collisionally ionized WHIM are ruled out at high confidence. We suggest that O VI gas is primarily associated with a photoionized CGM with $n_H \approx 10^{-5} \text{ cm}^{-3}$ and typical dimension $d \sim 300 \text{ kpc}$.

J. X. P. is partially supported by an NSF CAREER grant (AST-0548180). We acknowledge valuable comments and criticism of this work from B. Oppenheimer, G. Worseck, J. Tumlinson, and A. Fox.

REFERENCES

- Bahcall, J. N., & Spitzer, L. J. 1969, *ApJ*, 156, L63
 Barton, E. J., & Cooke, J. 2009, *AJ*, 138, 1817
 Bergeron, J. 1986, *A&A*, 155, L8
 Bergeron, J., & Boisse, P. 1991, *Advances in Space Research*, 11, 241
 Blanton, M. R., et al. 2003, *ApJ*, 594, 186
 Bond, N. A., Strauss, M. A., & Cen, R. 2010, *MNRAS*, 409, 156
 Bowen, D. V., Pettini, M., & Blades, J. C. 2002, *ApJ*, 580, 169
 Bregman, J. N., Miller, E. D., Athey, A. E., & Irwin, J. A. 2005, *ApJ*, 635, 1031
 Burles, S., & Tytler, D. 1996, *ApJ*, 460, 584
 Cen, R., & Fang, T. 2006, *ApJ*, 650, 573
 Cen, R., McDonald, P., Trac, H., & Loeb, A. 2009, *ArXiv e-prints*
 Cen, R., & Ostriker, J. P. 1999, *ApJ*, 514, 1
 —. 2006, *ApJ*, 650, 560
 Chen, H., Helsby, J. E., Gauthier, J., Sheckman, S. A., Thompson, I. B., & Tinker, J. L. 2010, *ApJ*, 714, 1521
 Chen, H., Lanzetta, K. M., Webb, J. K., & Barcons, X. 1998, *ApJ*, 498, 77
 Chen, H., & Mulchaey, J. S. 2009, *ApJ*, 701, 1219
 Chen, H.-W., Lanzetta, K. M., & Webb, J. K. 2001a, *ApJ*, 556, 158
 Chen, H.-W., Lanzetta, K. M., Webb, J. K., & Barcons, X. 2001b, *ApJ*, 559, 654
 Chen, H.-W., & Prochaska, J. X. 2000, *ApJ*, 543, L9
 Chen, H.-W., Prochaska, J. X., Weiner, B. J., Mulchaey, J. S., & Williger, G. M. 2005, *ApJ*, 629, L25
 Chen, H.-W., & Tinker, J. L. 2008, *ArXiv e-prints*, 801
 Cooksey, K. L., Prochaska, J. X., Chen, H.-W., Mulchaey, J. S., & Weiner, B. J. 2008, *ApJ*, 676, 262
 Cooksey, K. L., Thom, C., Prochaska, J. X., & Chen, H. 2010, *ApJ*, 708, 868
 Courteau, S., Dutton, A. A., van den Bosch, F. C., MacArthur, L. A., Dekel, A., McIntosh, D. H., & Dale, D. A. 2007, *ApJ*, 671, 203
 Danforth, C. W., & Shull, J. M. 2008, *ApJ*, 679, 194
 Danforth, C. W., Shull, J. M., Rosenberg, J. L., & Stocke, J. T. 2006, *ApJ*, 640, 716
 Davé, R., et al. 2001, *ApJ*, 552, 473
 Davé, R., Hernquist, L., Katz, N., & Weinberg, D. H. 1999, *ApJ*, 511, 521
 Davé, R., Oppenheimer, B. D., Katz, N., Kollmeier, J. A., & Weinberg, D. H. 2010, *MNRAS*, 408, 2051
 Dunkley, J., et al. 2009, *ApJS*, 180, 306
 Fox, A. J. 2011, *ApJ*, 730, 58
 Freeland, E., Cardoso, R. F., & Wilcots, E. 2008, *ApJ*, 685, 858
 Ganguly, R., Cen, R., Fang, T., & Sembach, K. 2008, *ArXiv e-prints*, 803
 Gnedin, N. Y., & Hui, L. 1998, *MNRAS*, 296, 44
 Gott, J. R., Choi, Y., Park, C., & Kim, J. 2009, *ApJ*, 695, L45
 Haardt, F., & Madau, P. 1996, *ApJ*, 461, 20
 Hibbard, J. E., van Gorkom, J. H., Rupen, M. P., & Schiminovich, D. 2001, in *Astronomical Society of the Pacific Conference Series*, Vol. 240, *Gas and Galaxy Evolution*, ed. J. E. Hibbard, M. Rupen, & J. H. van Gorkom, 657–+
 Lanzetta, K. M., Bowen, D. V., Tytler, D., & Webb, J. K. 1995, *ApJ*, 442, 538
 Lehner, N., Prochaska, J. X., Kobulnicky, H. A., Cooksey, K. L., Howk, J. C., Williger, G. M., & Cales, S. L. 2009, *ApJ*, 694, 734
 Lehner, N., Savage, B. D., Richter, P., Sembach, K. R., Tripp, T. M., & Wakker, B. P. 2007, *ApJ*, 658, 680
 Maller, A. H., & Bullock, J. S. 2004, *MNRAS*, 355, 694
 McDonald, P., et al. 2006, *ApJS*, 163, 80
 Miralda-Escudé, J., Cen, R., Ostriker, J. P., & Rauch, M. 1996, *ApJ*, 471, 582
 Mo, H. J., & Miralda-Escudé, J. 1996, *ApJ*, 469, 589
 Morris, S. L., & Jannuzi, B. T. 2006, *MNRAS*, 367, 1261
 Morris, S. L., Weymann, R. J., Dressler, A., McCarthy, P. J., Smith, B. A., Terrile, R. J., Giovanelli, R., & Irwin, M. 1993, *ApJ*, 419, 524
 Mulchaey, J. S., Davis, D. S., Mushotzky, R. F., & Burstein, D. 1996, *ApJ*, 456, 80
 Narayanan, A., Wakker, B. P., & Savage, B. D. 2009, *ApJ*, 703, 74
 Oppenheimer, B. D., & Davé, R. 2008, *MNRAS*, 387, 577
 —. 2009, *MNRAS*, 395, 1875
 Paschos, P., Jena, T., Tytler, D., Kirkman, D., & Norman, M. L. 2009, *MNRAS*, 399, 1934
 Penton, S. V., Stocke, J. T., & Shull, J. M. 2002, *ApJ*, 565, 720
 —. 2004, *ApJS*, 152, 29
 Prochaska, J. X., Chen, H.-W., Howk, J. C., Weiner, B. J., & Mulchaey, J. 2004, *ApJ*, 617, 718
 Prochaska, J. X., & Tumlinson, J. 2008, *ArXiv e-prints*
 Prochaska, J. X., Weiner, B. J., Chen, H., Cooksey, K. L., & Mulchaey, J. S. 2011, *apjsupp*
 Prochaska, J. X., Weiner, B. J., Chen, H.-W., & Mulchaey, J. S. 2006, *ApJ*, 643, 680
 Rauch, M. 1998, *ARA&A*, 36, 267
 Richter, P., Fang, T., & Bryan, G. L. 2006, *A&A*, 451, 767
 Rubin, K. H. R., Weiner, B. J., Koo, D. C., Martin, C. L., Prochaska, J. X., Coil, A. L., & Newman, J. A. 2010, *ApJ*, 719, 1503
 Savage, B. D., & de Boer, K. S. 1979, *ApJ*, 230, L77
 Savage, B. D., & Lehner, N. 2006, *ApJS*, 162, 134
 Savage, B. D., Lehner, N., Wakker, B. P., Sembach, K. R., & Tripp, T. M. 2005, *ApJ*, 626, 776
 Savage, B. D., & Sembach, K. R. 1991, *ApJ*, 379, 245
 Savage, B. D., et al. 2000, *ApJ*, 538, L27
 —. 2003, *ApJS*, 146, 125
 Sembach, K. R., Tripp, T. M., Savage, B. D., & Richter, P. 2004, *ApJS*, 155, 351
 Sembach, K. R., Wakker, B. P., Savage, B. D., & Richter, P. 2006, in *Astronomical Society of the Pacific Conference Series*, Vol. 348, *Astrophysics in the Far Ultraviolet: Five Years of Discovery with FUSE*, ed. G. Sonneborn, H. W. Moos, & B.-G. Andersson, 375–+

- Sembach, K. R., et al. 2003, *ApJS*, 146, 165
- Shone, A. M., Morris, S. L., Crighton, N., & Wilman, R. J. 2010, *MNRAS*, 402, 2520
- Spinrad, H., et al. 1993, *AJ*, 106, 1
- Spitzer, Jr., L. 1956, *ApJ*, 124, 20
- Stocke, J. T., Penton, S. V., Danforth, C. W., Shull, J. M., Tumlinson, J., & McLin, K. M. 2006, *ApJ*, 641, 217
- The SDSS Collaboration et al. 2010, *ArXiv e-prints*
- Thom, C., & Chen, H. 2008a, *ApJS*, 179, 37
- . 2008b, *ApJ*, 683, 22
- Tripp, T. M., Aracil, B., Bowen, D. V., & Jenkins, E. B. 2006, *ApJ*, 643, L77
- Tripp, T. M., Lu, L., & Savage, B. D. 1998, *ApJ*, 508, 200
- Tripp, T. M., Savage, B. D., & Jenkins, E. B. 2000, *ApJ*, 534, L1
- Tripp, T. M., Sembach, K. R., Bowen, D. V., Savage, B. D., Jenkins, E. B., Lehner, N., & Richter, P. 2008, *ApJS*, 177, 39
- Tumlinson, J., & Fang, T. 2005, *ApJ*, 623, L97
- Tumlinson, J., Shull, J. M., Giroux, M. L., & Stocke, J. T. 2005, *ApJ*, 620, 95
- Viel, M., Bolton, J. S., & Haehnelt, M. G. 2009, *MNRAS*, 399, L39
- Wakker, B. P., & Savage, B. D. 2009, *ApJS*, 182, 378
- Weiner, B. J., et al. 2009, *ApJ*, 692, 187
- Weymann, R. J., et al. 1998, *ApJ*, 506, 1
- Williger, G. M., Heap, S. R., Weymann, R. J., Davé, R., Ellingson, E., Carswell, R. F., Tripp, T. M., & Jenkins, E. B. 2006, *ApJ*, 636, 631
- Wilman, R. J., Morris, S. L., Jannuzi, B. T., Davé, R., & Shone, A. M. 2007, *MNRAS*, 375, 735
- Wise, J. H., Turk, M. J., Norman, M. L., & Abel, T. 2010, *ArXiv e-prints*
- York, D. G. 1974, *ApJ*, 193, L127
- Zheng, Z., Coil, A. L., & Zehavi, I. 2007, *ApJ*, 667, 760